

**Appendix C**  
Biological Reports



**Appendix C**  
**Table of Contents**

Results of Protocol Coastal California Gnatcatcher and Least Bell’s Vireo Surveys .....	C-1
Focused Survey Results for Sensitive Plant Species .....	C-24
Habitat Restoration Guidelines and Priorities.....	C-38
Summary of Focused Plan, Incidental and Protocol Survey Results .....	C-51
Focused Survey Results, Special-Status Plant Species .....	C-75
Least Bell’s Vireo and Coastal California Gnatcatcher Survey Results .....	C-105





GLENN LUKOS ASSOCIATES

Regulatory Services



July 26, 2010

Sandra Marquez  
U.S. Fish and Wildlife Service  
Carlsbad Field Office  
6010 Hidden Valley Road  
Carlsbad, California 92011

Lyann Comrack  
Nongame Wildlife Program  
California Department of Fish and Game  
1812 Ninth Street  
Sacramento, California 95811

**SUBJECT:** Results of Protocol Coastal California Gnatcatcher and Least Bell's Vireo Surveys for an Approximately 270-Acre Property Owned by the City of Whittier and Managed by the Puente Hills Landfill Native Habitat Preservation Authority, City of Whittier, Los Angeles County, California.

Dear Ms. Marquez:

This letter report documents the results of protocol presence/absence surveys conducted by Glenn Lukos Associates, Inc. (GLA) for the federally listed threatened coastal California gnatcatcher (*Poliophtila californica californica*) and the federally listed endangered least Bell's vireo (*Vireo bellii pusillus*). Focused surveys were conducted for a proposed oil extraction project within lands owned by the City of Whittier and managed by the Puente Hills Landfill Native Habitat Preservation Authority (Habitat Authority).

Surveys were conducted from April 23 through June 14, 2010 for the coastal California gnatcatcher, and from April 23 through July 7, 2010 for the least Bell's vireo. Focused surveys were conducted in all areas of potentially suitable habitat in accordance with U.S. Fish and Wildlife Service (USFWS) guidelines.

The least Bell's vireo was not detected within the survey area. The coastal California gnatcatcher was detected within the survey area, the results of which are discussed below.

**SURVEY AREA**

Focused surveys were conducted for a 270-acre survey area within lands owned by the City of Whittier and managed by the Habitat Authority [Exhibit 1 – Regional Map]. The survey area is located within Sections 22, 23, and 26, Township 2 South, Range 11 West [Exhibit 2 – Vicinity Map]. The Universal Transverse Mercator (UTM) coordinates approximately corresponding to

29 Orchard  
Telephone: (949) 837-0404

▪ Lake Forest

▪ California 92630-8300  
Facsimile: (949) 837-5834

Sandra Marquez  
U.S. Fish and Wildlife Service  
July 26, 2010  
Page 2

Lyann Comrack  
California Department of Fish and Game

the survey area are 407254 mE and 3759192 mN (Zone 11). The survey area is bordered by open space to the north, west, and east; residential development to the south and southwest, and Colima Road to the southeast. The topography of the survey area is generally characterized as high ridges bisected by two steep canyons, including La Cañada Verde and Arroyo Pescadero. A significant portion of the survey area has been disturbed in the past, in part by prior natural resource extraction activities. Portions of these areas have regenerated with native vegetation. Numerous dirt access roads and trails occur within the survey area, including actively maintained roads/trails and former roads that have become overgrown with vegetation. A ranger residence occurs within the southwest portion of the survey area. Portions of the survey area are accessed by the public for multiple purpose recreation; including the Arroyo Pescadero Trail and Deer Loop Trail, both of which are accessed from Colima Road to the southeast. Two native restoration sites occur within the southern portion of the survey area, one located between La Cañada Verde and Arroyo Pescadero, and the other at the Arroyo Pescadero trailhead.

Approximately 133.77 acres of the survey area support native vegetation communities, including coastal sage scrub (62.34 acres), chaparral (61.15 acres), and various riparian communities (10.28 acres). The majority of native upland scrub vegetation occurs in the northern portion of the survey area, on either side of La Cañada Verde Canyon, although a significant amount of native communities occur in the southeast portion of the study area. Coastal sage scrub areas are dominated by coastal sagebrush (*Artemisia californica*) and California buckwheat (*Eriogonum fasciculatum*), but also include other representative sage scrub species such as black sage (*Salvia mellifera*), white sage (*Salvia apiana*), and California brittlebush (*Encelia californica*). Chaparral areas are dominated by evergreen shrubs such as laurel sumac (*Malosma laurina*), lemonadeberry (*Rhus integrifolia*), coyote brush (*Baccharis pilularis*), and toyon (*Heteromeles arbutifolia*). Various riparian habitats occur within La Cañada Verde and Arroyo Pescadero, with the higher quality riparian vegetation occurring at the extreme northern end of the study area within La Cañada Verde. Dominant species at this location include arroyo willow (*Salix lasiolepis*), black willow (*Salix gooddingii*), and mule fat (*Baccharis salicifolia*). The remainder of La Cañada Verde contains a significant amount of non-native vegetation, including poison hemlock (*Conium maculatum*), but also contains occasional patches of Mexican elderberry (*Sambucus mexicanus*), and mule fat. Much of the overstory of both canyons consists of eucalyptus woodland. The portion of Arroyo Pescadero within the survey area consists mainly of eucalyptus woodland and a small patch of willows. The bottom of Arroyo Pescadero contains limited riparian vegetation, consisting of patches of elderberry and mule fat.

A smaller amount of the coastal sage scrub vegetation within the overall survey area (approximately 12.16 acres) is considered suitable gnatcatcher habitat to the extent that the habitat has a reasonable potential to support breeding pairs, including an area where a breeding pair was identified in the northern portion of the survey area. This includes patches of habitat in

Sandra Marquez  
U.S. Fish and Wildlife Service  
July 26, 2010  
Page 3

Lyann Comrack  
California Department of Fish and Game

the northern portion of the survey, and other patches in the southern portion of the site. Similarly, a limited amount of riparian habitat has the potential to support the least Bell's vireo (approximately 1.94 acres), consisting of a stand of southern willow scrub located within La Cañada Verde in the northern portion of the survey area [Exhibit 3 – Survey Area Map].

## **METHODOLOGY**

Protocol surveys for the coastal California gnatcatcher were performed in all suitable areas of coastal sage scrub, and to some extent areas of chaparral and mixed chaparral/sage scrub. Surveys were conducted in accordance with the 1997 USFWS guidelines, which stipulate that during the breeding season, six surveys shall be conducted in all areas of suitable habitat with at least seven days between site visits. The USFWS survey guidelines also stipulate that no more than 80 acres of suitable habitat shall be surveyed per biologist per day. The overall survey area contains approximately 60 acres of coastal sage scrub and 60 acres of chaparral, with the assumption that the survey area contains more than 80 acres but less than 160 acres of suitable habitat for the gnatcatcher. Therefore, the survey area was divided into two survey polygons requiring the equivalent of two “survey-days” per week (no more than 80 acres per day per biologist).

Protocol surveys for Polygon A were conducted on April 23, May 3, 13, and 24, and June 3 and 14, 2010. Surveys for Polygon B were conducted on April 26, May 3, 10, 17, and 26, and June 2, 2010. Surveys were conducted by Jeff Ahrens (TE 052159-3), Kevin Livergood (TE-172638-0), and David Moskovitz (TE-084606-1). The Habitat Authority's ecologist (Shannon Lucas) accompanied GLA biologists during the gnatcatcher surveys on April 23, and 26, and May 3, 10, 17, and 26. All surveys were conducted during the morning hours and were completed before 12:00 P.M. No surveys were conducted during extreme weather conditions (i.e., winds exceeding 15 miles per hour, rain, or temperatures in excess of 35°C). All areas of suitable habitat were surveyed on foot by walking slowly and methodically. Taped vocalizations and “pishing” sounds were utilized to elicit a response from gnatcatchers that might be present. Table 1 provides a summary of gnatcatcher survey dates.

Sandra Marquez  
 U.S. Fish and Wildlife Service  
 July 26, 2010  
 Page 4

Lyann Comrack  
 California Department of Fish and Game

**Table 1. Summary of Coastal California Gnatcatcher Survey Dates.**

Date	Polygon	Start Time	End Time	Permitted Surveyor	Temp °F (start/end)	Wind speed (mph)	Cloud Cover
4/23/2010	A	0715	1200	KL	50 / 67	0-2	clear
4/26/2010	B	0650	1200	JA	61 / 75	0-2	overcast
5/3/2010	A & B	0720	1130	JA/KL	57 / 77	0-2	clear
5/10/2010	B	0620	1100	JA	60 / 75	0-3	partly cloudy
5/13/2010	A	0625	1100	JA	62 / 74	0-2	partly cloudy
5/17/2010	B	0700	0945	KL	60 / 64	0-3	overcast
5/24/2010	A	0550	0920	JA	55 / 73	0-2	overcast
5/26/2010	B	0630	0930	KL	60 / 74	0-2	partly cloudy
6/2/2010	B	0630	0830	KL	60 / 62	0-1	overcast
6/3/2010	A	0630	1100	DM	62 / 66	0	overcast
6/14/2010	A	0610	1000	JA	59 / 81	0-1	Clear

KL – Kevin Livergood, JA – Jeff Ahrens, DM – David Moskovitz

Protocol surveys for the least Bell's vireo were conducted in areas of potentially suitable habitat, in accordance with the 1999 USFWS guidelines, which stipulate that a minimum of eight visits be conducted within areas of suitable habitat, with at least ten days between site visits. Biologists are to survey up to 50 hectares (approximately 120 acres) and no more than 3 linear kilometers (approximately 1.8 miles) per day, depending on site conditions (e.g., density and width of vegetation).

Protocol surveys were conducted April 23, May 3, 13 and 24, June 3, 14, and 24, and July 6, 2010. Surveys were conducted by David Moskovitz, Jeff Ahrens, Alisa Flint, and Kevin Livergood. All surveys were conducted between dawn and 11:00 a.m., in accordance with USFWS guidelines. All suitable areas were covered on foot by walking slowly and methodically through and adjacent to the riparian habitat. Birds were identified by call and sight, aided by the use of binoculars. No taped vocalizations were used to elicit response from vireos or any other species potentially present. No surveys were conducted during extreme weather conditions (i.e., winds exceeding 15 miles per hour, rain, or temperatures in excess of 35°C). Table 2 provides a summary of vireo survey dates.

Sandra Marquez  
 U.S. Fish and Wildlife Service  
 July 26, 2010  
 Page 5

Lyann Comrack  
 California Department of Fish and Game

**Table 2. Summary of Least Bell's Vireo Survey Dates.**

Date	Start Time	End Time	Surveying Biologist	Temp °F (start/end)	Wind speed (mph)	Cloud Cover
4/23/2010	0710	1045	DM	50 / 65	0-2	clear
5/3/2010	0700	1045	DM	57 / 66	0	clear
5/13/2010	0625	1100	JA	62 / 74	0-2	partly cloudy
5/24/2010	0550	0920	JA	55 / 73	0-2	overcast
6/3/2010	0630	1100	AF	62 / 66	0	overcast
6/14/2010	0610	1000	AF	59 / 81	0-1	clear
6/24/2010	0645	1000	AF	63 / 66	0-2	overcast
7/6/2010	0840	1030	KL	64 / 66	0	overcast

KL – Kevin Livergood, JA – Jeff Ahrens, DM – David Moskovitz, AF – Alisa Flint

## **RESULTS**

### ***Coastal California Gnatcatcher***

The coastal California gnatcatcher was observed in two locations within the survey area during protocol surveys, including one family group adjacent to La Cañada Verde in the northern portion of the survey area, and a single gnatcatcher adjacent to access road near the Worsham Landfill.

A single California gnatcatcher (sex unknown) vocalized once in response to tape playback on May 3, 2010. The response consisted of a single low-pitched mew and was barely audible. The gnatcatcher was utilizing coastal sage scrub vegetation on slopes to the west side of La Cañada Verde. The location was also in very close proximity to a blue-gray gnatcatcher (*Poliophtila caerulea*) pair that was located in the adjacent riparian habitat. On May 24, 2010, a male California gnatcatcher responded to tape playback in the same location where the single gnatcatcher was detected on May 3, 2010 (N. 38° 58' 43.563 latitude, W. 118° 0' 20.808 longitude). After observing the gnatcatcher foraging and vocalizing for approximately two to three minutes, the male then flew across the dirt access road into the riparian habitat where a female California gnatcatcher and two juveniles were also observed and briefly heard. The blue-gray gnatcatcher pair was also detected in very close proximity to the California gnatcatcher family group and briefly interacted with the group. On June 14, 2010, one California gnatcatcher was briefly observed in the same general location as the previous two detections. The bird was actively foraging and could only be identified by the underside of the retrice (tail) feathers. This bird was foraging in very close proximity to a blue-gray gnatcatcher family group and was presumed to be one member of the previously detected family group.

Sandra Marquez  
U.S. Fish and Wildlife Service  
July 26, 2010  
Page 6

Lyann Comrack  
California Department of Fish and Game

The habitat in which the California gnatcatcher family group was located consisted of a mixture of coastal sage scrub and chaparral plant species including California sagebrush, purple sage (*Salvia leucophylla*), white sage, Mexican elderberry, and laurel sumac. The riparian habitat in which the family group was observed in was comprised primarily of arroyo willow and mule fat, with some saltcedar (*Tamarix ramosissima*). The nest location of the pair could not be confirmed, though it was most likely located in the northern portion of the survey area. This pair was not detected during previous gnatcatcher surveys in 2005, 2008, or 2009.

The second gnatcatcher location consisted of a single male gnatcatcher that was observed on June 14, 2010 within scrub vegetation along the access road to the Worsham Landfill (N. 38° 59' 2.068 latitude, 118° 0' 48.445 longitude). The bird was detected in a location where LSA Associates observed a single gnatcatcher in 2005, though it is unclear whether it was the same bird. The habitat in which the California gnatcatcher was located consisted of thin strip of coastal sage scrub that included California sagebrush, purple sage, white sage, Mexican elderberry, and laurel sumac. Exhibit 3 depicts the locations of observed gnatcatchers.

#### ***Least Bell's Vireo***

The least Bell's vireo was not detected within the survey area during protocol surveys. Two other special-status riparian birds were detected during focused surveys, including the yellow-breasted chat (*Icteria virens*) and the yellow warbler (*Dendroica petechia brewsteri*). The yellow-breasted chat and one yellow warbler were detected in the northern portion of the survey area within La Cañada Verde. A second yellow warbler was detected north of the Arroyo Pescadero parking lot.




Sandra Marquez  
U.S. Fish and Wildlife Service  
July 26, 2010  
Page 7

Lyann Comrack  
California Department of Fish and Game

If you have any questions regarding the methodology or findings of this report, please contact David Moskovitz at (949) 837-0404, ext 42.

I certify that the information in this survey report and attached exhibits fully and accurately represents my work.


GLENN LUKOS ASSOCIATES, INC.

	TE-084606-1	July 26, 2010
David Moskovitz Biologist	Permit #	Date
	TE 052159-3	July 26, 2010
Jeff Ahrens Biologist	Permit #	Date
	TE-172638-0	July 26, 2010
Kevin Livergood Biologist	Permit #	Date

s:0953-02a.cagn\_lbv.rpt.c.doc



Adapted from USGS Santa Ana  
and Los Angeles Quadrangles



**CITY OF WHITTIER  
OIL EXPLORATION PROJECT**  
Regional Map


GLENN LUKOS ASSOCIATES



EXHIBIT 1



Adapted from USGS Whittier, CA quadrangle  
NORTH  
0 1,000 2,000 4,000  
Feet




**CITY OF WHITTIER  
OIL EXPLORATION PROJECT**  
Vicinity Map

GLENN LUKOS ASSOCIATES






EXHIBIT 2










**Legend**

-  CAGN Polygon A
-  CAGN Polygon B
-  LBV Suitable Habitat
-  CAGN Most Suitable Habitat
-  CAGN Occupied Habitat

**CAGN Locations**

-  Pair plus 2 juveniles
-  Single Adult



**CITY OF WHITTIER  
OIL EXPLORATION PROJECT**  
Survey Area Map



GLENN LUKOS ASSOCIATES 

Exhibit 3 

X:\0363-THE REST\0953-02OIL\953-2.GIS\953-2CAGN.mxd  
July 12, 2010



## FAUNAL COMPENDIUM

The faunal compendium lists species that were observed or detected by sign (e.g, tracks, scat, and burrows) within the Study Area. Non-native species are denoted by a ‘\*’. Sensitive species detected on site according to their status (e.g, breeding, wintering, rookery, etc.) as per the CDFG Special Animals List (July 2009) are denoted by ‘+’. Taxonomy and common names are taken from Pelham 2008 for butterflies, AOU (2009) for birds; CDFG (2008) for reptiles and amphibians; and CDFG (2008) for mammals.

### LEPIDOPTERA

#### HESPERIIDAE

*Erynnis funeralis*  
*Hylephila phyleus*  
*Atalopedes campestris*  
*Poanes melane*

#### PAPILIONIDAE

*Papilio zelicaon*  
*Papilio eurymedon*

#### PIERIDAE

*Pontia protodice*  
*Pieris rapae*  
*Anthocharis sara*  
*Colias philodice*  
*Colias eurytheme*

#### LYCAENIDAE

*Leptotes marina*  
*Cupido amyntula*  
*Hemiargus ceraunus*  
*Plebejus acmon*

#### NYMPHALIDAE

*Precis coenia*  
*Nymphalis antiopa*  
*Vanessa atalanta*  
*Vanessa cardui*  
*Vanessa anabella*  
*Vanessa virginiensis*  
*Liminitis lorquini*

### BUTTERFLIES

#### Skippers

funereal duskywing  
fiery skipper  
sachem  
umber skipper

#### Swallowtails

anise swallowtail  
pale swallowtail

#### Whites and Sulphurs

checkered white  
cabbage white  
Pacific orangetip  
clouded sulphur  
orange sulphur

#### Gossamer-Wing Butterflies

marine blue  
western tailed-blue  
Ceraunus blue  
acmon blue

#### Brush-Footed Butterflies

common buckeye  
mourning cloak  
red admiral  
painted lady  
west coast lady  
American lady  
Lorquin’s admiral

## REPTILIA

### PHRYNOSOMATIDAE

*Uta stansburiana*  
*Sceloporus occidentalis*

### COLUBRIDAE

*Pituophis catenifer*

### VIPERIDAE

*Crotalus oreganos or viridis*

## AVES

### ODONTOPHORIDAE

*Callipepla californica*

### CATHARTIDAE

*Cathartes aura*

### ACCIPITRIDAE

*Circus cyaneus*  
*Accipiter cooperii*  
*Buteo lineatus*  
*Buteo swainsoni*  
*Buteo jamaicensis*

### FALCONIDAE

*Falco sparverius*

### CHARADRIIDAE

*Charadrius vociferus*

### LARIDAE

*Larus delawarensis*  
*Larus occidentalis*  
*Larus californicus*

### COLUMBIDAE

- \* *Columba livia*
- Patagioenas fasciata*
- \* *Streptopelia decaocto*
- Zenaida macroura*

## REPTILES

### Phrynosomatid Lizards

common side-blotched lizard  
western fence lizard

### Colubrid Snakes

gopher snake

### Vipers

western rattlesnake

## BIRDS

### New World Quails

California quail

### New World Vultures

turkey vulture

### Hawks And Old World Vultures

northern harrier  
Cooper's hawk  
red-shouldered hawk  
Swainson's hawk  
red-tailed hawk

### Caracaras And Falcons

American kestrel

### Plovers And Relatives

killdeer

### Skuas, Gulls, Terns And Skimmers

ring-billed gull  
western gull  
California gull

### Pigeons And doves

rock pigeon  
band-tailed pigeon  
Eurasian collared-dove  
mourning dove

**CUCULIDAE**

*Geococcyx californianus*

**TYTONIDAE**

*Tyto alba*

**STRIGIDAE**

*Bubo virginianus*

**CAPRIMULGIDAE**

*Phalaenoptilus nuttallii*

**APODIDAE**

*Aeronautes saxatilis*

**TROCHILIDAE**

*Archilochus alexandri*

*Calypte anna*

*Selasphorus sasin*

**PICIDAE**

*Melanerpes formicivorus*

*Picoides nuttallii*

*Picoides pubescens*

*Colaptes auratus*

**TYRANNIDAE**

*Contopus cooperi*

*Contopus sordidulus*

*Empidonax difficilis*

*Sayornis nigricans*

*Sayornis saya*

*Myiarchus cinerascens*

*Tyrannus vociferans*

*Tyrannus verticalis*

**VIREONIDAE**

*Vireo huttoni*

*Vireo gilvus*

**CORVIDAE**

*Aphelocoma californica*

*Corvus brachyrhynchos*

*Corvus corax*

**Cuckoos, Roadrunners, And Anis**

greater roadrunner

**Barn Owls**

barn owl

**Typical Owls**

great horned owl

**GOATSUCKERS**

common poorwill

**Swifts**

white-throated swift

**Hummingbirds**

black-chinned hummingbird

Anna's hummingbird

Allen's hummingbird

**Woodpeckers And Allies**

acorn woodpecker

Nuttall's woodpecker

downy woodpecker

northern flicker

**Tyrant Flycatchers**

olive-sided flycatcher

western wood-pewee

Pacific-slope flycatcher

black phoebe

Say's phoebe

ash-throated flycatcher

Cassin's kingbird

western kingbird

**Vireos**

Hutton's vireo

warbling vireo

**Crows And Jays**

western scrub-jay

American crow

common raven

**HIRUNDINIDAE**

*Stelgidopteryx serripennis*  
*Petrochelidon pyrrhonota*

**AEGITHALIDAE**

*Psaltriparus minimus*

**TROGLODYTIDAE**

*Thryomanes bewickii*  
*Troglodytes aedon*

**REGULIDAE**

*Regulus calendula*

**SYLVIIDAE**

*Polioptila caerulea*  
+ *Polioptila californica californica*

**TURDIDAE**

*Sialia mexicana*  
*Catharus ustulatus*  
*Turdus migratorius*

**TIMALIIDAE**

*Chamaea fasciata*

**MIMIDAE**

*Mimus polyglottos*  
*Toxostoma redivivum*

**STURNIDAE**

\* *Sturnus vulgaris*

**BOMBYCILLIDAE**

*Bombycilla cedrorum*

**PTILOGONATIDAE**

*Phainopepla nitens*

**PARULIDAE**

*Vermivora celata*  
*Vermivora ruficapilla*  
+ *Dendroica petechia*  
*Dendroica coronata*  
*Dendroica nigrescens*  
*Geothlypis trichas*

**Swallows**

northern rough-winged swallow  
cliff swallow

**Long-Tailed Tits And Bushtits**

bushtit

**Wrens**

Bewick's wren  
house wren

**Kinglets**

ruby-crowned kinglet

**Old World Warblers And Gnatcatchers**

blue-gray gnatcatcher  
coastal California gnatcatcher

**Thrushes**

western bluebird  
Swainson's thrush  
American robin

**Babblers**

wrentit

**Mockingbirds And Thrashers**

northern mockingbird  
California thrasher

**Starlings And Allies**

European starling

**Waxwings**

cedar waxwing

**Silky-flycatchers**

phainopepla

**Wood Warblers And Relatives**

orange-crowned warbler  
Nashville warbler  
yellow warbler  
yellow-rumped warbler  
black-throated gray warbler  
common yellowthroat

*Wilsonia pusilla*  
 + *Icteria virens*

Wilson's warbler  
 yellow-breasted chat

**EMBERIZIDAE**

*Pipilo maculatus*  
*Pipilo crissalis*  
 + *Aimophila ruficeps*  
*Chondestes grammacus*  
*Passerculus sandwichensis*  
*Melospiza melodia*  
*Zonotrichia leucophrys*

**Emberizids**

spotted towhee  
 California towhee  
 rufous-crowned sparrow  
 lark sparrow  
 savannah sparrow  
 song sparrow  
 white-crowned sparrow

**CARDINALIDAE**

*Piranga ludoviciana*  
*Pheucticus melanocephalus*  
*Passerina caerulea*  
*Passerina amoena*

**Cardinals, Grosbeaks And Allies**

western tanager  
 black-headed grosbeak  
 blue grosbeak  
 lazuli bunting

**ICTERIDAE**

*Sturnella neglecta*  
*Euphagus cyanocephalus*  
*Molothrus ater*  
*Icterus cucullatus*  
*Icterus bullockii*

**Blackbirds**

western meadowlark  
 Brewer's blackbird  
 brown-headed cowbird  
 hooded oriole  
 Bullock's oriole

**FRINGILLIDAE**

*Carpodacus mexicanus*  
*Spinus psaltria*  
*Spinus tristis*

**Fringilline And Cardueline Finches and Allies**

house finch  
 lesser goldfinch  
 American goldfinch

**PASSERIDAE**

\* *Passer domesticus*

**Old World Sparrows**

house sparrow

**MAMMALIA**

**MAMMALS**

**DIDELPHIDAE**

\* *Didelphis virginiana*

**Opossums**

Virginia opossum

**LEPORIDAE**

*Sylvilagus audubonii*

**Rabbits And Hares**

desert (Audubon's) cottontail

**GEOMYIDAE**

*Thomomys bottae*

**Pocket Gophers**

Botta's pocket gopher

**MURIDAE**

*Neotoma fuscipes*

**SCIURIDAE**

*Sciurus griseus*

*Spermophilus beecheyi*

**CANIDAE**

\* *Canis familiaris*

*Canis latrans*

**PROCYONIDAE**

*Procyon lotor*

**MEPHITIDAE**

*Mephitis mephitis*

**FELIDAE**

\* *Felis catus*

*Lynx rufus*

**CERVIDAE**

*Odocoileus hemionus*

**Mice, Rats And Voles**

dusky-footed woodrat

**Squirrels, Chipmunks, And Marmots**

western gray squirrel

California ground squirrel

**Foxes, Wolves And Allies**

feral dog

coyote

**Raccoons And Allies**

raccoon

**Skunks**

striped skunk

**Cats**

feral cat

bobcat

**Deer, Elk And Allies**

mule deer



Mail to:  
 California Natural Diversity Database  
 Department of Fish and Game  
 1807 13<sup>th</sup> Street, Suite 202  
 Sacramento, CA 95811  
 Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov

For Office Use Only	
Source Code _____	Quad Code _____
Elm Code _____	Occ. No. _____
EO Index No. _____	Map Index No. _____

Date of Field Work (mm/dd/yyyy): 5/24/2010

### California Native Species Field Survey Form

Scientific Name: Poliophtilia California California

Common Name: Coastal California gnatcatcher

Species Found?  Yes  No If not, why?

Total No. Individuals 4 Subsequent Visit?  yes  no

Is this an existing NDDDB occurrence?  Yes, Occ. #  no  unk.

Collection? If yes: \_\_\_\_\_  
Number Museum / Herbarium

Reporter: Jeff Ahrens

Address: 29 Orchard, Lake Forest, CA 92630

E-mail Address: jahrens@wetlandpermitting.com

Phone: (949) 837-0404 ext 40.

**Plant Information**

Phenology: \_\_\_\_\_% vegetative \_\_\_\_\_% flowering \_\_\_\_\_% fruiting

**Animal Information**

2 # adults 2 # juveniles # larvae # egg masses # unknown

wintering  breeding  nesting  rookery  burrow site  other

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

County: Los Angeles Landowner / Mgr.: City of Whittier

Quad Name: Whittier Elevation: \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  GPS Make & Model \_\_\_\_\_

**DATUM:** NAD27  NAD83  WGS84  Horizontal Accuracy \_\_\_\_\_ meters/feet

Coordinate System: UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)

Coordinates: Lat 33° 58' 43.563", Long -118° 02' 0.808"

**Habitat Description (plants & animals) plant communities, dominants, associates, substrates/soils, aspects/slope:**  
**Animal Behavior (Describe observed behavior, such as territoriality, foraging, singing, calling, copulating, perching, roosting, etc., especially for avifauna):**

Coastal Sage Scrub / chaparral / riparian

Pair w/ 2 juveniles using willow riparian habitat immediately adjacent to Coastal Sage Scrub / chaparral.

Please fill out separate form for other rare taxa seen at this site.

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor

Immediate AND surrounding land use: Habitat Preserve

Visible disturbances: None

Threats: possible oil extraction activities

Comments: Family group detected one time during Protocol Surveys, one individual detected in same area on one occasion.

**Determination:** (check one or more, and fill in blanks)

Keyed (cite reference): \_\_\_\_\_

Compared with specimen housed at: \_\_\_\_\_

Compared with photo / drawing in: \_\_\_\_\_

By another person (name): \_\_\_\_\_

Other: \_\_\_\_\_

**Photographs:** (check one or more)

Plant / animal	<input type="checkbox"/>	Slide	<input type="checkbox"/>	Print	<input type="checkbox"/>	Digital	<input type="checkbox"/>
Habitat	<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>
Diagnostic feature	<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>

May we obtain duplicates at our expense? yes  no

Mail to:  
 California Natural Diversity Database  
 Department of Fish and Game  
 1807 13<sup>th</sup> Street, Suite 202  
 Sacramento, CA 95811  
 Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov

For Office Use Only	
Source Code _____	Quad Code _____
Elm Code _____	Occ. No. _____
EO Index No. _____	Map Index No. _____

Date of Field Work (mm/dd/yyyy): 6/14/2010

**California Native Species Field Survey Form**

**Scientific Name:** Poliophtilia californica californica

**Common Name:** Coastal California gnatcatcher

**Species Found?**  Yes  No If not, why? \_\_\_\_\_

Total No. Individuals 1 Subsequent Visit?  yes  no

Is this an existing NDDB occurrence?  yes, Occ. # \_\_\_\_\_  no  unk.

Collection? If yes: \_\_\_\_\_  
Number Museum / Herbarium

**Reporter:** Jeff Ahrens

**Address:** 29 Orchard, Lake Forest CA 92630

**E-mail Address:** jahrens@wetlandpermitting.com

**Phone:** (949) 837-0404 ext 40

Plant Information	Animal Information																				
Phenology: _____% vegetative _____% flowering _____% fruiting	<table border="0"> <tr> <td># adults</td> <td># juveniles</td> <td># larvae</td> <td># egg masses</td> <td># unknown</td> </tr> <tr> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> <tr> <td>wintering</td> <td>breeding</td> <td>nesting</td> <td>rookery</td> <td>burrow site</td> </tr> <tr> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> </table>	# adults	# juveniles	# larvae	# egg masses	# unknown	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	wintering	breeding	nesting	rookery	burrow site	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
# adults	# juveniles	# larvae	# egg masses	# unknown																	
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>																	
wintering	breeding	nesting	rookery	burrow site																	
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>																	

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

County: Los Angeles Landowner / Mgr.: City of Whittier

Quad Name: Whittier Elevation: \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  GPS Make & Model \_\_\_\_\_

**DATUM:** NAD27  NAD83  WGS84  Horizontal Accuracy \_\_\_\_\_ meters/feet

Coordinate System: UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)

Coordinates: Lat 33°59'2.068", Long -118°04'48.445"

**Habitat Description (plants & animals)** plant communities, dominants, associates, substrates/soils, aspects/slope:  
**Animal Behavior** (Describe observed behavior, such as territoriality, foraging, singing, calling, copulating, perching, roosting, etc., especially for avifauna):  
Bird detected in mixture of Coastal Sage Scrub & Chaparral

Please fill out separate form for other rare taxa seen at this site.

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor

Immediate AND surrounding land use: Habitat Preserve

Visible disturbances: Land fill

Threats: possible oil extraction activities

Comments: Individual detected only one time during Protocol Surveys,

<p><b>Determination:</b> (check one or more, and fill in blanks)</p> <input type="checkbox"/> Keyed (cite reference): _____ <input type="checkbox"/> Compared with specimen housed at: _____ <input type="checkbox"/> Compared with photo / drawing in: _____ <input type="checkbox"/> By another person (name): _____ <input type="checkbox"/> Other: _____	<p><b>Photographs:</b> (check one or more)</p> <table border="0"> <tr> <td>Plant / animal</td> <td><input type="checkbox"/></td> <td>Slide</td> <td><input type="checkbox"/></td> <td>Print</td> <td><input type="checkbox"/></td> <td>Digital</td> <td><input type="checkbox"/></td> </tr> <tr> <td>Habitat</td> <td><input type="checkbox"/></td> <td></td> <td><input type="checkbox"/></td> <td></td> <td><input type="checkbox"/></td> <td></td> <td><input type="checkbox"/></td> </tr> <tr> <td>Diagnostic feature</td> <td><input type="checkbox"/></td> <td></td> <td><input type="checkbox"/></td> <td></td> <td><input type="checkbox"/></td> <td></td> <td><input type="checkbox"/></td> </tr> </table> <p>May we obtain duplicates at our expense? yes <input type="checkbox"/> no <input type="checkbox"/></p>	Plant / animal	<input type="checkbox"/>	Slide	<input type="checkbox"/>	Print	<input type="checkbox"/>	Digital	<input type="checkbox"/>	Habitat	<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>	Diagnostic feature	<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>
Plant / animal	<input type="checkbox"/>	Slide	<input type="checkbox"/>	Print	<input type="checkbox"/>	Digital	<input type="checkbox"/>																		
Habitat	<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>																		
Diagnostic feature	<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>																		



Mail to:  
 California Natural Diversity Database  
 Department of Fish and Game  
 1807 13<sup>th</sup> Street, Suite 202  
 Sacramento, CA 95811  
 Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov

For Office Use Only	
Source Code _____	Quad Code _____
Elm Code _____	Occ. No. _____
EO Index No. _____	Map Index No. _____

Date of Field Work (mm/dd/yyyy): 4/23/2010

**Reset**

**California Native Species Field Survey Form**

**Send Form**

<b>Scientific Name:</b> <u>Icteria virens</u>	
<b>Common Name:</b> <u>Yellow-breasted chat</u>	
<b>Species Found?</b> <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No <small>If not, why?</small>	<b>Reporter:</b> <u>David Moskowitz</u>
Total No. Individuals <u>2</u> Subsequent Visit? <input type="checkbox"/> yes <input type="checkbox"/> no Is this an existing NDDDB occurrence? <input type="checkbox"/> no <input checked="" type="checkbox"/> unk. <small>Yes, Occ. #</small>	<b>Address:</b> <u>29 orchard, Lake forest, CA 92630</u>
Collection? If yes: _____ <small>Number Museum / Herbarium</small>	<b>E-mail Address:</b> <u>dmoskowitz@wetlandpermitting.com</u> <b>Phone:</b> <u>(449) 837-0404 ext 42</u>

**Plant Information**

Phenology: \_\_\_\_\_% vegetative \_\_\_\_\_% flowering \_\_\_\_\_% fruiting

**Animal Information**

2

# adults <input type="checkbox"/>	# juveniles <input type="checkbox"/>	# larvae <input type="checkbox"/>	# egg masses <input type="checkbox"/>	# unknown <input type="checkbox"/>
wintering	breeding	nesting	rookery	burrow site
other				

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

County: Los Angeles Landowner / Mgr.: City of Whittier

Quad Name: Whittier Elevation: \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian: H  M  S  Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian: H  M  S  GPS Make & Model \_\_\_\_\_

**DATUM:** NAD27  NAD83  WGS84  Horizontal Accuracy \_\_\_\_\_ meters/feet

**Coordinate System:** UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)

**Coordinates:** Lat 33°58.734, Long -118°00.336

**Habitat Description (plants & animals)** plant communities, dominants, associates, substrates/soils, aspects/slope:  
**Animal Behavior** (Describe observed behavior, such as territoriality, foraging, singing, calling, copulating, perching, roosting, etc., especially for avifauna):  
willow riparian / Eucalyptus  
Pair detected multiple times in willow riparian & adjacent Eucalyptus patch.

Please fill out separate form for other rare taxa seen at this site.

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor

Immediate AND surrounding land use: Habitat Preserve

Visible disturbances: None

Threats: Possible oil extraction activities

Comments:

**Determination:** (check one or more, and fill in blanks)

Keyed (cite reference): \_\_\_\_\_

Compared with specimen housed at: \_\_\_\_\_

Compared with photo / drawing in: \_\_\_\_\_

By another person (name): \_\_\_\_\_

Other: \_\_\_\_\_

**Photographs:** (check one or more)

Plant / animal	Slide <input type="checkbox"/>	Print <input type="checkbox"/>	Digital <input type="checkbox"/>
Habitat	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Diagnostic feature	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

May we obtain duplicates at our expense? yes  no

DFG/BDB/1747 Rev. 6/16/09

<p style="text-align: center;">Mail to:                  California Natural Diversity Database                  Department of Fish and Game                  1807 13<sup>th</sup> Street, Suite 202                  Sacramento, CA 95811                  Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov</p>	<p style="text-align: center;"><i>For Office Use Only</i></p> Source Code _____ Quad Code _____ Elm Code _____ Occ. No. _____ EO Index No. _____ Map Index No. _____																				
<p>Date of Field Work (mm/dd/yyyy): <u>5/13/10</u></p>																					
<input type="button" value="Reset"/>	<b>California Native Species Field Survey Form</b>	<input type="button" value="Send Form"/>																			
<p><b>Scientific Name:</b> <u>Dendroica petechia brewsteri</u></p>																					
<p><b>Common Name:</b> <u>Yellow warbler</u></p>																					
<p><b>Species Found?</b> <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No _____  <small>If not, why?</small></p> <p>Total No. Individuals <u>1</u> Subsequent Visit? <input type="checkbox"/> yes <input type="checkbox"/> no</p> <p>Is this an existing NDDDB occurrence? <input checked="" type="checkbox"/> no <input type="checkbox"/> unk.  <small>Yes, Occ. #</small></p> <p>Collection? If yes: _____  <small>Number Museum / Herbarium</small></p>	<p><b>Reporter:</b> <u>Jeff Ahrens</u></p> <p><b>Address:</b> <u>29 orchard, Lake Forest, CA 92630</u></p> <p><b>E-mail Address:</b> <u>jahrens@wetlandpermitting.com</u></p> <p><b>Phone:</b> <u>(949) 837-0664 ext 40</u></p>																				
<p><b>Plant Information</b></p> <p>Phenology: _____% vegetative _____% flowering _____% fruiting</p>	<p><b>Animal Information</b></p> <table style="width: 100%; text-align: center;"> <tr> <td># adults</td> <td># juveniles</td> <td># larvae</td> <td># egg masses</td> <td># unknown</td> </tr> <tr> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> <tr> <td>wintering</td> <td>breeding</td> <td>nesting</td> <td>rookery</td> <td>burrow site</td> </tr> <tr> <td></td> <td></td> <td></td> <td></td> <td>other</td> </tr> </table>	# adults	# juveniles	# larvae	# egg masses	# unknown	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	wintering	breeding	nesting	rookery	burrow site					other
# adults	# juveniles	# larvae	# egg masses	# unknown																	
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>																	
wintering	breeding	nesting	rookery	burrow site																	
				other																	
<p><b>Location Description (please attach map AND/OR fill out your choice of coordinates, below)</b></p> <p>County: <u>Los Angeles</u> Landowner / Mgr.: <u>City of Whittier</u></p> <p>Quad Name: <u>Whittier</u> Elevation: _____</p> <p>T _____ R _____ Sec _____, _____ 1/4 of _____ 1/4, Meridian: H <input type="checkbox"/> M <input type="checkbox"/> S <input type="checkbox"/> Source of Coordinates (GPS, topo. map &amp; type): _____</p> <p>T _____ R _____ Sec _____, _____ 1/4 of _____ 1/4, Meridian: H <input type="checkbox"/> M <input type="checkbox"/> S <input type="checkbox"/> GPS Make &amp; Model _____</p> <p><b>DATUM:</b> NAD27 <input type="checkbox"/> NAD83 <input type="checkbox"/> WGS84 <input type="checkbox"/> Horizontal Accuracy _____ meters/feet</p> <p>Coordinate System: UTM Zone 10 <input type="checkbox"/> UTM Zone 11 <input type="checkbox"/> OR Geographic (Latitude &amp; Longitude) <input type="checkbox"/></p> <p>Coordinates: <u>Lat 33° 58.497, Long -118.06.411</u></p>																					
<p><b>Habitat Description (plants &amp; animals) plant communities, dominants, associates, substrates/soils, aspects/slope:</b></p> <p><b>Animal Behavior</b> (Describe observed behavior, such as territoriality, foraging, singing, calling, copulating, perching, roosting, etc., especially for avifauna):</p> <p><u>Euc withdrawn</u></p> <p><u>Individual singing.</u></p> <p>Please fill out separate form for other rare taxa seen at this site.</p>																					
<p><b>Site Information</b> Overall site/occurrence quality/viability (site + population): <input type="checkbox"/> Excellent <input type="checkbox"/> Good <input checked="" type="checkbox"/> Fair <input type="checkbox"/> Poor</p> <p>Immediate AND surrounding land use: <u>Habitat Preserve</u></p> <p>Visible disturbances: <u>None</u></p> <p>Threats: <u>Potential oil extraction activities</u></p> <p>Comments:</p>																					
<p><b>Determination:</b> (check one or more, and fill in blanks)</p> <p><input type="checkbox"/> Keyed (cite reference): _____</p> <p><input type="checkbox"/> Compared with specimen housed at: _____</p> <p><input type="checkbox"/> Compared with photo / drawing in: _____</p> <p><input type="checkbox"/> By another person (name): _____</p> <p><input type="checkbox"/> Other: _____</p>	<p><b>Photographs:</b> (check one or more)</p> <table style="width: 100%; text-align: center;"> <tr> <td>Plant / animal</td> <td>Slide <input type="checkbox"/></td> <td>Print <input type="checkbox"/></td> <td>Digital <input type="checkbox"/></td> </tr> <tr> <td>Habitat</td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> <tr> <td>Diagnostic feature</td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> </table> <p>May we obtain duplicates at our expense? yes <input type="checkbox"/> no <input type="checkbox"/></p>	Plant / animal	Slide <input type="checkbox"/>	Print <input type="checkbox"/>	Digital <input type="checkbox"/>	Habitat	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	Diagnostic feature	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>								
Plant / animal	Slide <input type="checkbox"/>	Print <input type="checkbox"/>	Digital <input type="checkbox"/>																		
Habitat	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>																		
Diagnostic feature	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>																		



Mail to:  
 California Natural Diversity Database  
 Department of Fish and Game  
 1807 13<sup>th</sup> Street, Suite 202  
 Sacramento, CA 95811  
 Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov

For Office Use Only	
Source Code _____	Quad Code _____
Elm Code _____	Occ. No. _____
EO Index No. _____	Map Index No. _____

Date of Field Work (mm/dd/yyyy): 4/26/10

### California Native Species Field Survey Form

Scientific Name: Dendroica petechia brewsteri

Common Name: Yellow warbler

Species Found?  Yes  No If not, why?

Total No. Individuals 1 Subsequent Visit?  yes  no

Is this an existing NDDB occurrence?  no  unk. Yes, Occ. #

Collection? If yes: \_\_\_\_\_  
Number Museum / Herbarium

Reporter: Jeff Ahrens

Address: 29 Orchard, Lake Forest, CA 92630

E-mail Address: jahrens@wetlandpermitting.com

Phone: (499) 837-0404 ext 40

**Plant Information**

Phenology: \_\_\_\_\_% vegetative \_\_\_\_\_% flowering \_\_\_\_\_% fruiting

**Animal Information**

# adults	# juveniles	# larvae	# egg masses	# unknown
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
wintering	breeding	nesting	rookery	burrow site
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
other				

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

County: Los Angeles Landowner / Mgr.: City of Whittier

Quad Name: Whittier Elevation: \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  GPS Make & Model \_\_\_\_\_

**DATUM:** NAD27  NAD83  WGS84  Horizontal Accuracy \_\_\_\_\_ meters/feet

Coordinate System: UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)

Coordinates: Lat + 33° 58.021, Long - 118 00.073

**Habitat Description (plants & animals) plant communities, dominants, associates, substrates/soils, aspects/slope:**

**Animal Behavior** (Describe observed behavior, such as territoriality, foraging, singing, calling, copulating, perching, roosting, etc., especially for avifauna):

Eucalyptus patch.

Individual singing from near top of crown.

Please fill out separate form for other rare taxa seen at this site.

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor

Immediate AND surrounding land use: Habitat Preserve

Visible disturbances: None

Threats: Possible oil extraction activities

Comments:

**Determination:** (check one or more, and fill in blanks)

Keyed (cite reference): \_\_\_\_\_

Compared with specimen housed at: \_\_\_\_\_

Compared with photo / drawing in: \_\_\_\_\_

By another person (name): \_\_\_\_\_

Other: \_\_\_\_\_

**Photographs:** (check one or more)

Plant / animal	<input type="checkbox"/>	Slide	<input type="checkbox"/>	Print	<input type="checkbox"/>	Digital	<input type="checkbox"/>
Habitat	<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>
Diagnostic feature	<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>

May we obtain duplicates at our expense? yes  no

Mail to:  
 California Natural Diversity Database  
 Department of Fish and Game  
 1807 13<sup>th</sup> Street, Suite 202  
 Sacramento, CA 95811  
 Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov

For Office Use Only	
Source Code _____	Quad Code _____
Elm Code _____	Occ. No. _____
EO Index No. _____	Map Index No. _____

Date of Field Work (mm/dd/yyyy): 5/24/2010

**California Native Species Field Survey Form**

<b>Scientific Name:</b> <u>Aimophila ruficeps canescens</u>	
<b>Common Name:</b> <u>Southern California rufous-crowned sparrow</u>	
<b>Species Found?</b> <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No _____ <small>If not, why?</small>	<b>Reporter:</b> <u>Jeff Ahrens</u>
Total No. Individuals <u>1</u> Subsequent Visit? <input type="checkbox"/> yes <input type="checkbox"/> no Is this an existing NDDDB occurrence? <input checked="" type="checkbox"/> no <input type="checkbox"/> unk. <small>Yes, Occ. #</small>	<b>Address:</b> <u>29 Orchard, Lake Forest, CA 92630</u>
Collection? If yes: _____ <small>Number Museum / Herbarium</small>	<b>E-mail Address:</b> <u>jahrens@wetlandpermitting.com</u> <b>Phone:</b> <u>(949) 837-0444 ext 46</u>

**Plant Information**

Phenology: \_\_\_\_\_% vegetative \_\_\_\_\_% flowering \_\_\_\_\_% fruiting

**Animal Information**

# adults	# juveniles	# larvae	# egg masses	# unknown
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
wintering	breeding	nesting	rookery	burrow site
other				

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

County: Los Angeles Landowner / Mgr.: City of Whittier

Quad Name: Whittier Elevation: \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  GPS Make & Model \_\_\_\_\_

**DATUM:** NAD27  NAD83  WGS84  Horizontal Accuracy \_\_\_\_\_ meters/feet

Coordinate System: UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)

Coordinates: Lat 33.58687, long -118.00443

**Habitat Description (plants & animals)** plant communities, dominants, associates, substrates/soils, aspects/slope:  
**Animal Behavior** (Describe observed behavior, such as territoriality, foraging, singing, calling, copulating, perching, roosting, etc., especially for avifauna):  
Coastal Sage scrub / chaparral

Please fill out separate form for other rare taxa seen at this site.

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor

Immediate AND surrounding land use: Habitat preserve

Visible disturbances: None

Threats: Possible oil extraction activities

Comments:

**Determination:** (check one or more, and fill in blanks)

Keyed (cite reference): \_\_\_\_\_

Compared with specimen housed at: \_\_\_\_\_

Compared with photo / drawing in: \_\_\_\_\_

By another person (name): \_\_\_\_\_

Other: \_\_\_\_\_

**Photographs:** (check one or more)

Plant / animal	<input type="checkbox"/>	Slide	<input type="checkbox"/>	Print	<input type="checkbox"/>	Digital	<input type="checkbox"/>
Habitat	<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>
Diagnostic feature	<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>		<input type="checkbox"/>

May we obtain duplicates at our expense? yes  no

DFG/BDB/1747 Rev. 6/16/09



Mail to:  
 California Natural Diversity Database  
 Department of Fish and Game  
 1807 13<sup>th</sup> Street, Suite 202  
 Sacramento, CA 95811  
 Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov

For Office Use Only	
Source Code _____	Quad Code _____
Elm Code _____	Occ. No. _____
EO Index No. _____	Map Index No. _____

Date of Field Work (mm/dd/yyyy): 5/26/2016

**Reset**

**California Native Species Field Survey Form**

**Send Form**

**Scientific Name:** Aimophila ruficeps canescens

**Common Name:** southern California rufous-crowned sparrow

**Species Found?**  Yes  No \_\_\_\_\_ If not, why? \_\_\_\_\_

Total No. Individuals 1 Subsequent Visit?  yes  no

Is this an existing NDDDB occurrence? \_\_\_\_\_  no  unk.  
Yes, Occ. # \_\_\_\_\_

Collection? If yes: \_\_\_\_\_  
Number \_\_\_\_\_ Museum / Herbarium \_\_\_\_\_

**Reporter:** Kevin Livergood

**Address:** 29 Orchard, Lake Forest, CA 92630

**E-mail Address:** klivergood@wetlandpermitting.com

**Phone:** (949) 837-0404 ext 36

**Plant Information**

Phenology: \_\_\_\_\_% vegetative \_\_\_\_\_% flowering \_\_\_\_\_% fruiting

**Animal Information**

1

# adults	# juveniles	# larvae	# egg masses	# unknown
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
wintering	breeding	nesting	rookery	burrow site
other				

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

County: Los Angeles Landowner / Mgr.: City of Whittier

Quad Name: Whittier Elevation: \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  GPS Make & Model \_\_\_\_\_

**DATUM:** NAD27  NAD83  WGS84  Horizontal Accuracy \_\_\_\_\_ meters/feet

Coordinate System: UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)

Coordinates: Lat 33.967869, Long -117.998397

**Habitat Description (plants & animals)** plant communities, dominants, associates, substrates/soils, aspects/slope:  
**Animal Behavior** (Describe observed behavior, such as territoriality, foraging, singing, calling, copulating, perching, roosting, etc., especially for avifauna):  
Coastal sage scrub & chaparral

Please fill out separate form for other rare taxa seen at this site.

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor

Immediate AND surrounding land use: Habitat Preserve

Visible disturbances: None

Threats: Possible oil extraction activities

Comments:

**Determination:** (check one or more, and fill in blanks)

Keyed (cite reference): \_\_\_\_\_

Compared with specimen housed at: \_\_\_\_\_

Compared with photo / drawing in: \_\_\_\_\_

By another person (name): \_\_\_\_\_

Other: \_\_\_\_\_

**Photographs:** (check one or more)

Slide	Print	Digital
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Plant / animal	<input type="checkbox"/>	<input type="checkbox"/>
Habitat	<input type="checkbox"/>	<input type="checkbox"/>
Diagnostic feature	<input type="checkbox"/>	<input type="checkbox"/>

May we obtain duplicates at our expense? yes  no

DFG/BDB/1747 Rev. 6/16/09

LSA

LSA ASSOCIATES, INC.  
1500 IOWA AVENUE, SUITE 200  
RIVERSIDE, CALIFORNIA 92507

951.781.9310 TEL  
951.781.4277 FAX

OTHER OFFICES:  
IRVINE  
PT. RICHMOND  
SAN LUIS OBISPO  
PALM SPRINGS  
FORT COLLINS  
BERKELEY  
ROCKLIN  
SOUTH SAN FRANCISCO  
CARLSBAD  
FRESNO

July 19, 2010

Ms. Andrea Gullo  
Puente Hills Landfill Native Habitat Preservation Authority  
7702 Washington Avenue, Suite C  
Whittier, California 90602

Subject: Focused Survey Results for Sensitive Plant Species, City of Whittier Oil Exploration  
(LSA Project No. PUE0901)

Dear Ms. Gullo:

This letter report documents the results of focused plant surveys conducted by LSA Associates, Inc. (LSA) in 2008, 2009, and 2010 for the proposed oil exploration activities within lands managed by the Puente Hills Landfill Native Habitat Preservation Authority (Habitat Authority), owned by and located in the City of Whittier, Los Angeles County, California (Figure 1; all figures are attached).

Sensitive plant species were determined to be absent from the survey areas. There were no incidental observations of western spadefoot (*Spea hammondi*) or cactus wren (*Campylorhynchus brunneicapillus*) during the surveys.

## BACKGROUND

Areas surveyed in 2008 totaled approximately 113 acres under investigation for potential oil exploration activities. The area surveyed in 2009 totaled approximately 209 acres and overlapped with much of the land surveyed in 2008 (Figure 2). Two areas, totaling approximately 40 acres, were surveyed in 2010. These two areas are distinct from those surveyed in 2008 and 2009. The survey areas are located within Sections 22, 23, 25, and 26 of Township 2 South, Range 11 West, as shown on the United States Geological Survey (USGS) 7.5-minute *Whittier* and *La Habra, California* quadrangles (Figure 1).

The survey areas are characterized by portions of Arroyo Pescadero and La Cañada Verde drainages, adjacent hillsides, and access roads. Elevations range from approximately 300 to 1,000 feet above sea level. Vegetation types within the survey areas primarily include coastal sage scrub, riparian scrub, nonnative grassland, ruderal vegetation, eucalyptus woodland, ornamental vegetation, and previously disturbed communities. Soil types mapped in the survey areas (Appendix B of LSA 2007) include the Hanford Association (0 to 5% slopes), Perkins-Rincon Association (0 to 15% slopes), and Altamont-Diablo Association (30 to 60% slopes, eroded).

7/19/2010 (R:\PUE0901\Botany\2008-10 Plant Survey Results\_Fin.doc)

PLANNING | ENVIRONMENTAL SCIENCES | DESIGN



LSA ASSOCIATES, INC.

Prior to conducting focused surveys, LSA biologists reviewed appropriate literature to determine whether sensitive plant species have been detected on or near the survey areas in the past. The literature review included the results of previous surveys of the project site (LSA 2006 and 2009) and a published checklist of plants of the Whittier Hills (Ljubenkov and Ross 2002), as well as the California Natural Diversity Database (CNDDDB; California Department of Fish and Game 2008) and the California Native Plant Society's (CNPS) Online Inventory (California Native Plant Society 2008). The CNDDDB query included the *La Habra* and *Whittier, California* quadrangles, and the CNPS query included a 9-quad search of the surrounding quadrangles. LSA also reviewed the Resource Management Plan (LSA 2007) prepared for the Habitat Authority in 2007, including the Sensitive Species Table in Appendix I, to further refine which sensitive plant species might be present in the survey areas.

Based on the literature review, no sensitive plants were found to have historic locations within the survey areas. However, nearby records for Plummer's mariposa lily (*Calochortus plummerae*) and Robinson's peppergrass (*Lepidium virginicum* var. *robinsonii*), both on CNPS List 1B, and Catalina mariposa lily (*Calochortus catalinae*), a CNPS List 4 species, combined with the presence of potentially suitable habitat, suggested that there was potential for these species to occur within the survey areas. Also, Southern California black walnut (*Juglans californica*), a CNPS List 4 species, was observed on the site during the 2009 survey. While the timing and methods of surveys focused on these four target species, all vascular plant species observed in the survey areas were identified to the degree necessary to determine sensitivity status.

## METHODS

LSA biologists surveyed the respective survey areas for each year according to the following schedule:

LSA Biologists	Date	Time
Jim Harrison, Dan Rosie	April 10, 2008	7:00 a.m. to 4:30 p.m.
Dan Rosie, Jodi Ross	April 18, 2008	6:00 a.m. to 12:30 p.m.
Jim Harrison, Matthew Willis	June 4, 2008	6:30 a.m. to 3:30 p.m.
Jim Harrison, Dan Rosie	June 5, 2008	8:00 a.m. to 12:00 p.m.
Dan Rosie, Robert Steers	April 2, 2009	10:15 a.m. to 6:00 p.m.
Dan Rosie, Robert Steers	April 3, 2009	10:00 a.m. to 12:15 p.m.
Sarah Barrera, Robert Steers	June 5, 2009	11:00 a.m. to 3:30 p.m.
Sarah Barrera, Robert Steers	June 9, 2009	9:00 a.m. to 12:00 p.m.
Stan Spencer, Jodi Ross	April 27, 2010	10:30 a.m. to 2:00 p.m.
Stan Spencer, Jodi Ross	April 28, 2010	11:30 a.m. to 1:20 p.m.
Stan Spencer, Jodi Ross	June 10, 2010	10:00 a.m. to 1:00 p.m.

LSA ASSOCIATES, INC.

Botanical surveys were conducted in accordance with the current CNPS Botanical Survey Guidelines (California Native Plant Society 2001). In each year, the first site visit was conducted in April to observe plants that mature in early spring, and a later visit was conducted in June to observe plants that mature during late spring.

The surveys were conducted by walking transects throughout the survey areas. Transect widths varied from 10 to about 100 feet and depended on visibility and habitat quality. Although the surveys were conducted during the expected flowering seasons of the target species in order to facilitate detection of the plants, transects were walked slowly enough that the target species could have been detected even in a preflowering or postflowering state. Steep slopes inaccessible by foot were surveyed using binoculars.

Precipitation in the City of Whittier was 9.7 inches from September of 2007 to May 2008 and 8.4 inches from September of 2008 to May 2009 (National Climate Data Center 2009). Average precipitation for the City of Whittier is 14.05 inches from September through May, based on 59 years of data (Western Regional Climate Center 2009). Therefore, precipitation was below average in both of these survey years. However, the majority of rainfall occurred between November and February in both years (data not shown), which led to widespread germination of native annual plants and bolting of perennial geophytes such as blue dicks (*Dichelostemma capitatum*) and blue-eyed grass (*Sisyrinchium bellum*). Furthermore, in 2009 Habitat Authority ecologist Shannon Lucas confirmed that Plummer's mariposa lily was blooming at a nearby site during one of the survey visits (pers. com. June 4, 2009). Thus, it was concluded that these were adequate years and sampling dates to detect target plant species. Precipitation in the general site vicinity from September 2009 to May 2010 was above average (University of California 2010).

Attached Table A contains a cumulative list of plant species identified during the 2008, 2009, and 2010 surveys.

## RESULTS AND DISCUSSION

No sensitive plant species were detected during the surveys. A stand of about 30 individuals of various ages of Southern California black walnut was found in the drainage that is parallel to and east of Catalina Avenue (Figure 2). This stand is a component of the riparian vegetation that occurs in the drainage, and is disturbed, with eucalyptus trees (*Eucalyptus* sp.) predominating. Southern California black walnut is on the CNPS 4 List. CNPS List 4 is only a "watch list." Species on this list are not generally considered sensitive and do not appear on CNPS or CNDDB searches by USGS quads. This species has no State or Federal status but it is included in the Resource Management Plan for the preserve.

Historically, the survey areas have been heavily disturbed and much of the survey areas consist of nonnative vegetation. Patches of intact coastal sage scrub and other habitat potentially suitable for sensitive species do occur within the survey areas. However, the herbaceous component of these patches is dominated by nonnative species. Other portions of the survey areas that appeared relatively uninvaded were not found to contain sensitive plants. The combination of historic disturbance and a high abundance of nonnative species likely preclude the existence of sensitive plant species in the survey areas.

LSA ASSOCIATES, INC.


## CONCLUSIONS

Based on the results of the focused surveys, it is the conclusion of LSA that sensitive plant species do not occur within the areas surveyed.

If you have any questions or require additional information, please feel free to call me at (951) 781-9310.

Sincerely,

LSA ASSOCIATES, INC.

A handwritten signature in blue ink, appearing to read "Stanley C. Spencer".

Stanley C. Spencer, Ph.D.  
Senior Biologist

Attachments:   References Cited  
                  Table A: Plant Species Observed  
                  Figure 1: Project Location  
                  Figure 2: 2008, 2009, and 2010 Survey Areas and Results

LSA ASSOCIATES, INC.

## REFERENCES

- California Department of Fish and Game, Natural Heritage Division, Natural Diversity Database. 2008. RareFind Version 3.1.0. Records search executed April 8, 2008, covering the USGS 7.5-minute series topographic map, La Habra and Whittier, California quadrangles. Sacramento, California: The Resources Agency. Commercial version dated February 2, 2008.
- California Native Plant Society. 2001. Botanical Survey Guidelines. Revised June 2, 2001.
- . 2008. Inventory of Rare and Endangered Plants (online edition, v7-08b). California Native Plant Society. Sacramento, CA. Accessed on April 8, 2008. [<http://www.cnps.org/inventory>].
- Ljubenkov, J.A.S., and T.S. Ross. 2002. An Annotated Checklist of the Vascular Plants of the Whittier Hills, Los Angeles County, California. *Crossosoma* 27(1).
- LSA Associates, Inc. 2006. Botanical Survey Report 2005. Prepared for the Puente Hills Landfill Native Habitat Preservation Authority. January 13, 2006.
- LSA Associates, Inc. 2007. Resource Management Plan. Prepared for the Puente Hills Landfill Native Habitat Preservation Authority. July 2007.
- LSA Associates, Inc. 2009. Focused Survey Results, Special-Status Plant Species, City of Whittier Oil Exploration (LSA Project No. PUE0901). Prepared for the Puente Hills Landfill Native Habitat Preservation Authority. August 4, 2009.
- National Climatic Data Center. 2009. Annual Climatological Summary for Station: WHITTIER CITY YD FC106C, for 2007, 2008, and 2009. Website: <http://www.ncdc.noaa.gov/oa/climate/stationlocator.html> (Accessed on July 23, 2009).
- Western Regional Climate Center. 2009. Long Term Weather Summary for Station: WHITTIER CITY YD FC106C, Period of Record: 1/ 1/1949 to 12/31/2008. Website: <http://wrcc.dri.edu> (Accessed on July 23, 2009).
- University of California. 2010. UC IPM Online. California Weather Data for Pomona.A Station. Website: <http://www.ipm.ucdavis.edu/calludt.cgi/WXDESCRIPTION?STN=POMONA.A> (Accessed on July 7, 2010).

LSA ASSOCIATES, INC.

**TABLE A**  
**VASCULAR PLANT SPECIES OBSERVED**

The following vascular plant species were observed in the survey areas by various biologists during the course of on-site surveys in 2008, 2009, and 2010.

**Table A: Vascular Plant Species Observed**

Scientific Name	Common Name
<b>MAGNOLIOPHYTA: MAGNOLIOPSIDA</b>	<b>DICOT FLOWERING PLANTS</b>
<b>Anacardiaceae</b>	<b>Sumac family</b>
<i>Malosma laurina</i>	Laurel sumac
<i>Rhus integrifolia</i>	Lemonade berry
<i>Rhus ovata</i>	Sugar bush
<i>Schinus molle</i> (nonnative species)	Peruvian pepper tree
<i>Schinus terebinthifolius</i> (nonnative species)	Brazilian pepper tree
<i>Toxicodendron diversilobum</i>	Poison oak
<b>Apiaceae</b>	<b>Carrot family</b>
<i>Conium maculatum</i> (nonnative species)	Poison hemlock
<i>Daucus pusillus</i>	American wild carrot
<i>Foeniculum vulgare</i> (nonnative species)	Fennel
<b>Apocynaceae</b>	<b>Dogbane family</b>
<i>Vinca major</i> (nonnative species)	Blue periwinkle
<b>Asclepiadaceae</b>	<b>Milkweed family</b>
<i>Asclepias californica</i>	California milkweed
<b>Asteraceae</b>	<b>Sunflower family</b>
<i>Ambrosia acanthicarpa</i>	Annual bur-sage
<i>Ambrosia psilostachya</i>	Western ragweed
<i>Artemisia californica</i>	California sagebrush
<i>Artemisia douglasiana</i>	Mugwort
<i>Baccharis emoryi</i>	Emory's baccharis
<i>Baccharis pilularis</i>	Coyote brush
<i>Baccharis salicifolia</i>	Mule fat
<i>Carduus pycnocephalus</i> (nonnative species)	Italian Thistle
<i>Centaurea melitensis</i> (nonnative species)	Tocalote
<i>Cirsium vulgare</i> (nonnative species)	Bull thistle
<i>Corethrogyne filaginifolia</i>	California aster
<i>Cotula australis</i> (nonnative species)	Australian brass-buttons

LSA ASSOCIATES, INC.

**Table A: Vascular Plant Species Observed**

Scientific Name	Common Name
<i>Deinandra fasciculata</i>	Fascicled tarweed
<i>Encelia californica</i>	California encelia
<i>Eriophyllum confertiflorum</i>	Golden yarrow
<i>Gutierrezia californica</i>	California matchweed
<i>Hedynois cretica</i> (nonnative species)	Crete weed
<i>Helianthus annuus</i>	Common sunflower
<i>Heterotheca grandiflora</i>	Telegraph weed
<i>Isocoma menziesii</i>	Goldenbush
<i>Lactuca serriola</i> (nonnative species)	Prickly lettuce
<i>Logfia filaginoides</i>	California cottonrose
<i>Malacothrix saxatilis</i>	Cliff malacothrix
<i>Picris echioides</i> (nonnative species)	Bristly ox-tongue
<i>Pseudognaphalium biolettii</i>	Two-color rabbit-tobacco
<i>Pseudognaphalium californicum</i>	California rabbit-tobacco
<i>Pseudognaphalium luteoalbum</i> (nonnative species)	Jersey cudweed
<i>Pseudognaphalium microcephalum</i>	San Diego rabbit-tobacco
<i>Pseudognaphalium stramineum</i>	Cottonbatting plant
<i>Senecio vulgaris</i> (nonnative species)	Common groundsel
<i>Silybum marianum</i> (nonnative species)	Milk thistle
<i>Sonchus asper</i> (nonnative species)	Prickly sow thistle
<i>Sonchus oleraceus</i> (nonnative species)	Common sow thistle
<i>Stephanomeria virgata</i>	Tall wreath-plant
<i>Taraxacum officinale</i> (nonnative species)	Common dandelion
<i>Xanthium strumarium</i>	Rough cocklebur
<b>Bignoniaceae</b>	<b>Bignonia family</b>
<i>Jacaranda mimosifolia</i> (nonnative species)	Jacaranda
<b>Brassicaceae</b>	<b>Mustard family</b>
<i>Brassica nigra</i> (nonnative species)	Black mustard
<i>Hirschfeldia incana</i> (nonnative species)	Shortpod mustard
<i>Raphanus sativus</i> (nonnative species)	Wild radish
<i>Sisymbrium erysimoides</i> (nonnative species)	Mediterranean rocket
<i>Sisymbrium irio</i> (nonnative species)	London rocket
<b>Cactaceae</b>	<b>Cactus family</b>
<i>Opuntia littoralis</i>	Coastal pricklypear
<b>Caprifoliaceae</b>	<b>Honeysuckle family</b>
<i>Sambucus mexicana</i>	Blue elderberry

LSA ASSOCIATES, INC.

**Table A: Vascular Plant Species Observed**

Scientific Name	Common Name
<b>Caryophyllaceae</b>	<b>Pink family</b>
<i>Stellaria media</i> (nonnative species)	Common chickweed
<b>Chenopodiaceae</b>	<b>Saltbush family</b>
<i>Atriplex semibaccata</i> (nonnative species)	Australian saltbush
<i>Chenopodium album</i> (nonnative species)	Lamb's quarters
<i>Chenopodium berlandieri</i>	Pitseed goosefoot
<i>Chenopodium murale</i> (nonnative species)	Nettleleaf goosefoot
<i>Salsola tragus</i> (nonnative species)	Russian thistle
<b>Convolvulaceae</b>	<b>Morning-glory family</b>
<i>Calystegia macrostegia</i>	Morning-glory
<i>Convolvulus arvensis</i> (nonnative species)	Field bindweed
<b>Crassulaceae</b>	<b>Stonecrop family</b>
<i>Crassula comata</i>	Sand pigmy-stonecrop
<b>Cucurbitaceae</b>	<b>Gourd family</b>
<i>Cucurbita foetidissima</i>	Calabazilla
<i>Marah macrocarpus</i>	Cucamonga manroot
<b>Euphorbiaceae</b>	<b>Spurge family</b>
<i>Chamaesyce albomarginata</i>	Rattlesnake weed
<i>Chamaesyce maculata</i> (nonnative species)	Spotted spurge
<i>Ricinus communis</i> (nonnative species)	Castor bean
<b>Fabaceae</b>	<b>Pea family</b>
<i>Acacia cyclops</i> (nonnative species)	Coastal wattle
<i>Acacia longifolia</i>	Sydney golden wattle
<i>Lotus salsuginosus</i>	Coastal lotus
<i>Lotus scoparius</i>	Deerweed
<i>Lupinus microcarpus</i>	Chick lupine
<i>Lupinus succulentus</i>	Arroyo lupine
<i>Medicago polymorpha</i> (nonnative species)	Bur-clover
<i>Melilotus indicus</i> (nonnative species)	Annual yellow sweetclover
<i>Vicia villosa</i> (nonnative species)	Winter vetch
<b>Fagaceae</b>	<b>Beech family</b>
<i>Quercus agrifolia</i>	Coast live oak
<b>Geraniaceae</b>	<b>Geranium family</b>
<i>Erodium brachycarpum</i> or <i>botrys</i> (nonnative species)	Erodium
<i>Erodium cicutarium</i> (nonnative species)	Redstem stork's bill
<i>Erodium moschatum</i> (nonnative species)	Musky stork's bill

LSA ASSOCIATES, INC.

**Table A: Vascular Plant Species Observed**

Scientific Name	Common Name
<b>Grossulariaceae</b>	<b>Gooseberry family</b>
<i>Ribes speciosum</i>	Fuchsiaflower gooseberry
<b>Hydrophyllaceae</b>	<b>Waterleaf family</b>
<i>Emmenanthe penduliflora</i>	Whispering bells
<i>Eucrypta chrysanthemifolia</i>	Common eucrypta
<i>Phacelia cicutaria</i>	Caterpillar phacelia
<i>Phacelia distans</i>	Distant phacelia
<i>Phacelia minor</i>	Wild Canterbury bells
<i>Phacelia parryi</i>	Parry's phacelia
<i>Phacelia ramosissima</i>	Branching phacelia
<i>Phacelia tanacetifolia</i>	Tansy phacelia
<i>Pholistoma auritum</i>	Blue fiesta flower
<b>Juglandaceae</b>	<b>Walnut family</b>
<i>Juglans californica</i>	Southern California black walnut
<b>Lamiaceae</b>	<b>Mint family</b>
<i>Marrubium vulgare</i> (nonnative species)	Horehound
<i>Salvia apiana</i>	White sage
<i>Salvia leucophylla</i>	Purple sage
<i>Salvia leucophylla</i> X <i>apiana</i>	White/purple sage hybrid
<i>Salvia mellifera</i>	Black sage
<b>Lauraceae</b>	<b>Laurel family</b>
<i>Persea americana</i> (nonnative species)	Avocado
<b>Malvaceae</b>	<b>Mallow family</b>
<i>Malacothamnus fasciculatus</i>	Chaparral mallow
<i>Malva parviflora</i> (nonnative species)	Cheeseweed
<i>Malva sylvestris</i> (nonnative species)	High mallow
<b>Myrtaceae</b>	<b>Myrtle family</b>
<i>Eucalyptus</i> sp. (nonnative species)	Eucalyptus
<b>Nyctaginaceae</b>	<b>Four-o'clock family</b>
<i>Mirabilis laevis</i>	Wishbone bush
<b>Oleaceae</b>	<b>Olive family</b>
<i>Fraxinus velutina</i>	Velvet ash
<i>Ligustrum lucidum</i> (nonnative species)	Glossy privet
<b>Onagraceae</b>	<b>Evening primrose family</b>
<i>Camissonia californica</i>	Mustard-like evening primrose
<i>Clarkia botaiae</i>	Botta's clarkia



LSA ASSOCIATES, INC.

**Table A: Vascular Plant Species Observed**

Scientific Name	Common Name
<b>Oxalidaceae</b>	<b>Oxalis family</b>
<i>Oxalis pes-caprae</i> (nonnative species)	Bermuda buttercup
<b>Platanaceae</b>	<b>Sycamore family</b>
<i>Platanus racemosa</i>	Western sycamore
<b>Polygonaceae</b>	<b>Buckwheat family</b>
<i>Eriogonum fasciculatum</i>	California buckwheat
<i>Polygonum aviculare</i> (nonnative species)	Common knotweed
<i>Rumex crispus</i> (nonnative species)	Curly dock
<b>Portulacaceae</b>	<b>Purslane family</b>
<i>Claytonia perfoliata</i>	Miner's lettuce
<b>Primulaceae</b>	<b>Primrose family</b>
<i>Anagallis arvensis</i> (nonnative species)	Scarlet pimpernel
<b>Punicaceae</b>	<b>Pomegranate Family</b>
<i>Punica granatum</i> (nonnative species)	Pomegranate
<b>Rosaceae</b>	<b>Rose family</b>
<i>Eriobotrya japonica</i>	Loquat
<i>Heteromeles arbutifolia</i>	Toyon
<i>Prunus ilicifolia</i>	Hollyleaf cherry
<b>Rubiaceae</b>	<b>Madder family</b>
<i>Galium angustifolium</i> ssp. <i>angustifolium</i>	Narrow-leaved bedstraw
<i>Galium aparine</i>	Goose grass
<b>Salicaceae</b>	<b>Willow family</b>
<i>Salix exigua</i>	Narrowleaf willow
<i>Salix gooddingii</i>	Goodding's willow
<i>Salix laevigata</i>	Red willow
<i>Salix lasiolepis</i>	Arroyo willow
<b>Scrophulariaceae</b>	<b>Figwort family</b>
<i>Keckiella cordifolia</i>	Red bush penstemon
<i>Mimulus aurantiacus</i>	Red bush monkey-flower
<i>Scrophularia californica</i>	Coast figwort
<i>Verbascum virgatum</i> (nonnative species)	Wand Mullein
<b>Solanaceae</b>	<b>Nightshade family</b>
<i>Nicotiana glauca</i> (nonnative species)	Tree tobacco
<i>Solanum americanum</i>	American black nightshade
<i>Solanum douglasii</i>	Greenspot nightshade
<b>Tamaricaceae</b>	<b>Tamarisk family</b>
<i>Tamarix ramosissima</i> (nonnative species)	Mediterranean tamarisk

LSA ASSOCIATES, INC.

**Table A: Vascular Plant Species Observed**

Scientific Name	Common Name
<b>Tropaeolaceae</b>	<b>Nasturtium family</b>
<i>Tropaeolum majus</i> (nonnative species)	Garden nasturtium
<b>Ulmaceae</b>	<b>Elm family</b>
<i>Ulmus parvifolia</i> (nonnative species)	Chinese elm
<i>Ulmus pumila</i> (nonnative species)	Siberian elm
<b>Urticaceae</b>	<b>Nettle Family</b>
<i>Parietaria hespera</i>	Rillita pellitory
<i>Urtica dioica</i>	Stinging nettle
<i>Urtica urens</i> (nonnative species)	Dwarf nettle
<b>Verbenaceae</b>	<b>Vervain family</b>
<i>Verbena lasiostachys</i>	Western verbena
<b>Zygophyllaceae</b>	<b>Caltrop family</b>
<i>Tribulus terrestris</i> (nonnative species)	Puncture vine
<b>MAGNOLIOPHYTA: LILIOPSIDA</b>	<b>MONOCOT FLOWERING PLANTS</b>
<b>Arecaceae</b>	<b>Palm family</b>
<i>Washingtonia filifera</i>	California fan palm
<b>Iridaceae</b>	<b>Iris family</b>
<i>Sisyrinchium bellum</i>	Blue-eyed grass
<b>Liliaceae</b>	<b>Lily family</b>
<i>Dichelostemma capitatum</i>	Blue dicks
<i>Yucca gloriosa</i> (nonnative species)	Spanish dagger
<b>Poaceae</b>	<b>Grass family</b>
<i>Avena barbata</i> (nonnative species)	Slender wild oat
<i>Avena fatua</i> (nonnative species)	Wild oat
<i>Bromus diandrus</i> (nonnative species)	Ripgut brome
<i>Bromus hordeaceus</i> (nonnative species)	Soft chess
<i>Bromus madritensis</i> ssp. <i>rubens</i> (nonnative species)	Red brome
<i>Cortaderia jubata</i> (nonnative species)	Andean pampas grass, jubatagrass
<i>Cynodon dactylon</i> (nonnative species)	Bermuda grass
<i>Hordeum murinum</i> (nonnative species)	Foxtail barley
<i>Lamarckia aurea</i> (nonnative species)	Goldentop
<i>Leymus condensatus</i>	Giant wildrye
<i>Leymus triticoides</i>	Beardless wildrye
<i>Lolium multiflorum</i> (nonnative species)	Italian ryegrass
<i>Nassella lepida</i>	Foothill needlegrass
<i>Nassella pulchra</i>	Purple needlegrass
<i>Pennisetum setaceum</i> (nonnative species)	African fountain grass

LSA ASSOCIATES, INC.

**Table A: Vascular Plant Species Observed**

<b>Scientific Name</b>	<b>Common Name</b>
<i>Piptatherum miliaceum</i> (nonnative species)	Smilo grass
<i>Schismus barbatus</i> (nonnative species)	Common Mediterranean grass
<i>Vulpia myuros</i> (nonnative species)	Rat-tail fescue







FIGURE 1

LSA



- 2010 Survey Area (approx. 40 acres)
- 2009 Survey Area (approx. 209 acres)
- 2008 Survey Area (approx. 111 acres)


Puente Hills Landfill Native Habitat Preservation Authority  
City of Whittier Oil Exploration

Regional and Project Location


SOURCE: USGS 7.5' Quad: Whittier (81), CA; La Habra (81), CA; Thomas Bros., 2009

I:\Pue0901\Reports\Bio\fig1\_RegLoc.mxd (07/07/10)





LSA



- 2010 Survey Area (approx. 40 acres)
- 2009 Survey Area (approx. 209 acres)
- 2008 Survey Area (approx. 111 acres)
- Southern California Black Walnut (*Juglans californica*)

FIGURE 2

Puente Hills Landfill Native Habitat Preservation Authority  
City of Whittier Oil Exploration

2008, 2009, and 2010  
Survey Areas and Results

SOURCE: AirPhotoUSA, 2008, Thomas Bros., 2009

I:\Pue0901\Reports\Bio\fig2\_SurveyAreas.mxd (07/07/10)

## HABITAT RESTORATION GUIDELINES AND PRIORITIES

The purpose of this Habitat Restoration Plan (Plan) is to provide guidance on restoring degraded and disturbed habitats throughout the Habitat Authority property. While the Plan provides a great deal of technical information on existing conditions in the Preserve and on restoration methods, it is programmatic in nature and accomplishes the following:

- Identifies the range of conditions that exist in the potential restoration areas, specifically soil characteristics and weed composition;
- Provides restoration criteria and a priority evaluation on restoring the degraded and disturbed habitats;
- Provides information on the most effective restoration methods currently known and their associated costs;
- Provides basic data and recommendations prescribing restoration methods for each type of potential restoration area;
- Provides guidelines for preparing more detailed, site-specific plans that will maximize the success and minimize the cost of individual restoration efforts; and
- Provides guidance for approving future mitigation projects in the Preserve.

Specific plans for individual restoration sites should be developed on a case-by-case basis, with consideration of the information and guidelines provided in this Plan as well as new information that is developed through adaptive management.

This Plan is organized by the analyses of existing conditions (e.g., soil and weeds), restoration criteria and priority, restoration application, restoration techniques, performance standards and monitoring, and planting and seeding palettes.

This Plan considers all of the baseline resource and cultural resource data to make sure that the tenets of Ecosystem Management are incorporated. The Plan utilizes restoration criteria on which to base the restoration priorities as well as a master list of techniques and the situations for which they are appropriate. The restoration areas are evaluated for site conditions, and recommendations of the specific restoration techniques are prescribed for each type of restoration area.

### Approach

This Plan was prepared with three primary concepts in mind: Ecosystem Management, Adaptive Management, and Ecological Successional Model.

**Ecosystem Management.** Ecosystem Management integrates scientific knowledge of ecological relationships within a complex sociopolitical and values framework toward the general goal of protecting native ecosystem integrity over the long term.

The following are Ecosystem Management goals:

- Maintain viable populations of all native species in situ;
- Represent, within protected areas, all native ecosystem types across their natural range of variation;
- Maintain evolutionary and ecological processes (e.g., disturbance regimes, hydrological processes, nutrient cycles);
- Manage over a period of time long enough to maintain the evolutionary potential of species and ecosystems; and
- Accommodate human use and occupancy within these constraints.

**Adaptive Management.** Adaptive Management incorporates regular monitoring to evaluate the implemented Plan. Adaptive Management allows for continual adjustments to improve upon the current Plan. It is expected that this Plan will be used as a guide and that as more restoration is implemented in the Preserve, improvements will be made from each restoration success and failure.



**Ecological Successional Model.** The Ecological Successional Model mimics the successional process that occurs in nature following a disturbance. In nature, fast-growing plant species quickly recolonize the disturbed areas. These fast-growing species are well suited for competing against the heavily invasive alien species such as mustard, annual grasses, and thistle. In addition, these early seral species help prepare the soil by colonizing mycorrhizae and fixing nitrogen for the slower-developing perennials. By the time the vegetation reaches the climax plant community, most of the early successional species have dropped out of the plant community. However, these early successional species are lying dormant in the soil as seed, ready to germinate following the next disturbance. Plant communities are continuously in a state of change, constantly progressing towards a climax state, and are always being disturbed by natural and human forces. By basing the restoration primarily on seeded species, the specific site conditions will determine the actual climax plant community. These conditions and their effects on the ultimate community cannot always be known with certainty. In contrast, a climax restoration model attempts to mimic the climax plant community. This type of restoration leaves out the early successional species, primarily relying on container plants to provide the instant climax plant community. This model also assumes that the restoration “designer” knows what the climax community should be including its species composition.



## Soil

An understanding of soil and vegetation associations is key to determining appropriate habitat restoration. To start, LSA determined whether any of the soil associations were more likely to support exotic weeds. Table A-O shows the distribution of weedy areas across soil associations in relation to native vegetation. Table A-P shows that generally, exotic weeds are likely to be found in all soil associations from clay soils on gentle slopes to sandy loam soils on steep slopes.

**Table A-O: Soil Associations Acreage in Relation to Native Vegetation and Weed Distribution**

Soil Association	Soil Association Total Acres	Acres of Native Vegetation (%)	Acres of Weeds (%)
San Andreas-San Benito 30–70 percent slope	1,266	862 (68%)	404 (32%)
Hanford	618	360 (58%)	258 (42%)
Mocho-Sorrento	16	12 (75%)	4 (25%)
Perkins-Ricon	374	224 (60%)	150 (40%)
Altamont-Diablo 9–30 percent slope	341	238 (70%)	103 (30%)
Altamont-Diablo 30–50 percent slope	1,175	804 (68%)	371 (32%)

**Table A-P: General Relationships of Exotic Species**

Weed Community	Soil Characteristics		Aspect
	Texture	Calcareous (Lime Detected)	
<i>Brassica nigra/Centaurea melitensis</i>	Sandy Loam	No Lime	East to South to West
<i>Brassica nigra</i> /Nonnative grass	Clay Loam to Loam	Preference	All
<i>Brassica nigra/Silybum marianum</i>	Clay Loam	No Lime	East to South to West
<i>Erodium cicutarium</i> /Nonnative grass	Clay Loam	Preference	All
<i>Eucalyptus glauca</i>	Clay to Clay Loam	No Lime	All
<i>Foeniculum vulgare</i>	Clay to Clay Loam	No Lime	All
<i>Hirschfeldia incana/Centaurea melitensis</i>	Clay	Preference	West to Southeast
<i>Nicotiana glauca/Brassica nigra</i>	Clay Loam	Preference	South to Southwest
Nonnative grass/ <i>Brassica nigra</i>	Clay Loam, Clay to Loam	Preference	All



Weed Community	Soil Characteristics		Aspect
	Texture	Calcareous (Lime Detected)	
Nonnative grass/ <i>Centaurea melitensis</i>	Clay Loam	Preference	Southeast to Southwest
Nonnative grass/ <i>Erodium cicutarium</i>	Clay	No Lime	All
Nonnative grass/ <i>Eucalyptus glauca</i>	Clay	No Lime	All
Nonnative grass/ <i>Hirschfeldia incana</i>	Clay Loam to Clay	No Lime	All
Nonnative grass/ <i>Phalaris aquatica</i>	Clay	No Lime	North to Southeast
Nonnative grass/ <i>Pichris echioides</i>	Clay	No Lime	Northwest to East
Nonnative grass/ <i>Raphanus sativus</i>	Clay	No Lime	All
<i>Phalaris aquatica</i> /Nonnative grass	Clay	No Preference	Northwest to Northeast
<i>Raphanus sativus</i> / <i>Brassica nigra</i>	Clay to Clay Loam	No Lime	All
<i>Ricinus communis</i> / <i>Silybum marianum</i>	Loam	Preference	Southeast to West
<i>Schinus terebenthifolius</i> / <i>Brassica nigra</i>	Clay Loam	Preference	South to Southeast

The analyses from the Exotic Plant Species section (Appendix G) show the general relationships between soil, aspect, and weed species. These conclusions are based on limited soil tests.

Table A-Q shows the general relationship of some of the dominant native communities based on the limited soil testing conducted for this study. These general relationships can be used as a basis for developing the most appropriate native habitat for restoration in the Preserve. However, it should be stressed that the results are based on sample test locations over the entire Preserve. A more comprehensive sampling regime at specific locations for several key soil characteristics, such as lime, texture, and soil shrink-swell characteristics would provide more insight to guide appropriate habitat restoration.

**Table A-Q: Specific Relationships of Native Communities Based upon Limited Soil Tests**

Plant Community	Soil Characteristics		Aspect
	Texture	Calcareous (Lime Detected)	
Black Sage Scrub	Loam to Clay Loam	No Preference	East to West
Chaparral	Loam to Clay Loam	No Preference	North to Northwest
Coyote Brush Scrub	Clay	No Preference	Northwest to Southeast
Elderberry Woodland	Clay Loam	No Lime	North to West
Nassella Grassland	Clay to Clay Loam	No Lime	No Preference
Oak Woodland	Clay Loam to Loam	No Lime	North
Purple Sage Scrub	Clay Loam	Preference	Southeast to Southwest
Sagebrush Scrub	Sandy Loam to Clay	Low Preference	No Preference
Sagebrush/Buckwheat Scrub	Sandy Loam to Clay	No Lime	Southeast to Southwest
Walnut Woodland	Clay	Preference	Northeast to West

Based upon the results of LSA's analysis, which indicates that particular habitats prefer certain soil types, further soil investigation should be required during the development of a specific plan for each identified weed polygon. At a minimum, the soil should be mapped within each polygon to determine the overall type of soil: clay, clay loam, or loams. If the study is conducted during summer or early fall, then soil cracks should be noted to establish the shrink-swell capacity of the soils. Additionally, pooled soil samples from similar soil textures across the site should be collected, and tests for lime and available phosphorous should be performed. After these soil analyses establish texture and limited chemistry, then geomorphic position, slope, and aspect will contribute to determining an appropriate habitat for restoration based upon descriptions and analyses in the preceding sections.

### **Restoration Criteria and Priority Ranking**


Restoration criteria and priority ranking were developed with input from the Habitat Authority when all the data were collected and analyzed and results were discussed. The criteria and priorities will be analyzed for and applied to the previously identified weed polygons. It is important to note that weeds are scattered throughout the Preserve and not only limited to the areas mapped by BonTerra; however, the largest and highest concentration of weeds are found in these areas and will be the most useful for restoration planning purposes.

Habitat restoration/priorities were originally derived based on a concept of individual "management areas" (Whittier, Hacienda Heights, and La Habra Heights) throughout the Preserve (Figure A-7). However current management efforts are based on a Preserve-wide assessment. Therefore, the originization of the priorities by management unit provided herein is primarily for general information and does not prescribe actual management priorities.

Another factor affecting restoration priorities is the annual restoration budget. It will be important to maximize the restoration effort and cost-effectiveness to provide the most ecologically meaningful restoration.


**Priority Calculating Method.** Restoration priorities were developed using a number of factors including average slope category; polygon size; proximity to trails/roads; proximity to existing restoration efforts; whether it is positioned on a ridge top above natives; the presence of targeted highly invasive species and whether the targeted invasive species are the top two dominant species; and wildlife connectivity. Each category was given a priority value based upon criteria developed with input from the Habitat Authority. Although each priority value is somewhat subjective, weighting is based on the relative degree of difficulty for restoration and habitat value in an effort to maximize the amount of habitat restored within the Habitat Authority's budget. It is important to note that this analysis does not include fire or rare-plant data because they were not available at the time of this analysis. The rankings from each of the categories were added together, resulting in a priority ranking for the overall Preserve. The management areas were further divided into restoration planning units by watershed. Each of restoration planning units is referenced with the names called out on the USGS map. All unnamed restoration units are designated with a letter referencing the management area within the same watershed and a number. For example, H3 refers to the third canyon in the









Prepared By: L S A

FIGURE A-7



-  HACIENDA HEIGHTS MANAGEMENT AREA
-  LA HABRA HEIGHTS MANAGEMENT AREA
-  WHITTIER MANAGEMENT AREA



Puente Hills Landfill  
Native Habitat Preservation Authority

Resource Management Plan

Habitat Restoration Plan  
Management Areas

SOURCE: Aerial-EagleAerial (2003)  
F:\PUE430\GIS\Maps\Draft RMP\Appendices\FigA-7\_ManagementAreas.mxd (03/05/2007)



Hacienda Heights restoration unit. The restoration units were then ranked throughout the Preserve. Because the Preserve manages the land according to city/community ownership, the ranking of restoration units over the whole Preserve are further ranked by management area (Whittier, Hacienda Heights, and La Habra Heights). The restoration priority factors are described below.

**Slope.** In general, it is easier and less expensive to restore land with gentle slopes than land with steep slopes. The steeper areas are more difficult to access with equipment and personnel; tend to be more erosive; and, in extreme cases, can present a hazardous working condition. The percent slope was calculated for the weed polygons within the Preserve. The slope was broken into four categories: 0–20 percent, 20–40 percent, 40–60 percent, and 60–85 percent). Each weed polygon was designated the slope category with the most area for that polygon. Since some of the potential restoration areas are on very steep terrain, such as in Turnbull Canyon, these areas were given a low priority and ranked 2. The more gentle areas were ranked 40. The slope categories and priority values are shown in Table A-R below.

**Table A-R: Percent Slope Categories and Priority Values**

Percent Slope (%)	Priority Value
0–20	40
20–40	36
40–60	20
60–85	2

**Size.** The size of the weed polygons is generally related to a cost efficiency factor. The larger the area, the more cost-effective it will be to restore it. The largest weed polygons were designated a priority value of 10, and the smallest weed polygons were designated a priority value of 1. The weed polygon size categories and priority values are shown in Table A-S below.

**Table A-S: Weed Polygon Size Categories and Priority Values**

Weed Polygon Size	Priority Value
25–50 acres	10
10–25 acres	8
5–10 acres	6
1–5 acres	2
< 1 acre	1

**Proximity to Roads and Trails.** Site access by equipment and personnel is important when evaluating a restoration site. Site access was determined by proximity to existing roads or trails. The roads and trails were buffered at 10 feet, 50 feet, 100 feet, 500 feet, 1,000 feet, and 5,000 feet. The weed polygons were classified by the closest proximity category to the road or trail. Table A-T shows the priority-valued designated for each proximity classification.

**Table A-T: Proximity to Roads and Trails**

Proximity to Roads and Trails (feet)	Priority Value
< 10	10
10–50	9
50–100	8
101–500	5
501–1,000	3
1,000–5,000	2
> 5,000	1

**Proximity to Existing Restoration.** There are a number of restoration efforts that are planned or are currently underway in the Preserve. In order to help protect the integrity of these young restoration sites from composition from surrounding weeds, higher priority was given to those weed polygons in close proximity to existing or planned restoration sites. Also, the areas near existing restoration sites usually have well-traveled access and are nearby existing staging areas. Table A-U shows the priority values for proximity of existing restoration.

**Table A-U: Proximity to Existing Restoration Efforts**

Proximity to Existing Restoration (feet)	Priority Value
< 500	10
501–1,000	8
1,001–2,000	6
> 2,000	3

**Exotics' Position on Ridge Tops.** In areas where exotics are positioned at the highest elevations, natural conversion to native plant communities is the most difficult. These areas do not have a continuous source of native seeds as they would if positioned downhill of native plant communities. In addition, these exotics will continue to spread seed downhill into native plant communities. The weed polygons that are positioned on ridge tops are designated a priority value of 10, and the other weed polygons are designated a priority value of 4. Table A-V shows the priority value for the ridge top position.

**Table A-V: Exotics Positioned on Ridge Tops**

Exotics Positioned on Ridge Tops	Priority Value
Yes	10
No	4

**Highly Invasive Species.** There are some exotic species that are more invasive than others. The most highly invasive exotic weeds are identified and rated by California Invasive Plant Council (CalIPC). In addition, the Preserve has provided input on weeds that seem to be spreading in the Preserve. The most invasive of weeds should be a top priority to slow and stop their spread. If one or more of these species was present, the highest priority value was designated for that weed polygon. In addition, the amount of area these highly invasive weeds occupy is an important factor in their rate of spread and eradication. To account for this, weed polygons where the dominant and second most dominant weeds were invasive with a rating greater than 5 had a multiplier applied as follows. For weed polygons where the dominant weed was a species greater than 5, a 1.5 multiplier was applied. For weed polygons where the second dominant weed was a species greater than 5, a 1.2 multiplier was applied. The three numbers, including highly invasive weed species, most dominant invasive weed with a value greater than 5 (with multiplier), and second dominant highly invasive weed with a value greater than 5 (with multiplier), were added to the total. Table A-W shows a list of the most highly invasive weeds and their designated priority value.

**Table A-W: Highly Invasive Species and Priority Value**

Highly Invasive Exotic Species Present		CAL-IPC	Value
Scientific Name	Common Name		
<i>Schinus molle</i>	Peruvian pepper	Limited	3
<i>Nonnative Grasses</i>	NNG	Moderate	3
<i>Brassica nigra</i>	black mustard	Moderate	4
<i>Bromus diandrus</i>	ripgut brome	Moderate	4
<i>Robinia pseudoacacia</i>	black locust	Limited	5
<i>Nicotiana glauca</i>	tree tobacco	Moderate	6
<i>Carduus pycnocephalus</i>	Italian thistle	Moderate	10
<i>Cirsium arvense</i>	Canada thistle	Moderate	10
<i>Cirsium vulgare</i>	bull thistle	Moderate	10
<i>Cortaderia selloan</i>	pampas grass	High	10
<i>Foeniculum vulgare</i>	Fennel	High	10
<i>Myoporum laetum</i>	Myoporum	Moderate	10
<i>Pennisetum setaceum</i>	fountain grass	Moderate	10
<i>Phalaris aquatica</i>	harding grass	Moderate	10
<i>Ricinus communis</i>	castor bean	Limited	10
<i>Schinus terebinthifolius</i>	Brazilian pepper	Limited	10
<i>Silybum marianum</i>	milk thistle	Limited	10
<i>Conium maculatum</i>	poison hemlock	Moderate	10
<i>Eucalyptus sp.</i>	Eucalyptus	Limited to Moderate	10
<i>Acacia sp.</i>	Acacia	Limited	8
<i>Washingtonia robusta</i>	Mexican fan palm	Moderate	6

<b>Highly Invasive Exotic Species Present</b>		<b>CAL-IPC</b>	<b>Value</b>
<b>Scientific Name</b>	<b>Common Name</b>		
Dominance 1	> 5 Invasive Value	Multiply by 1.5	
Dominance 2	> 5 Invasive Value	Multiply by 1.2	

**Wildlife Connectivity.** The Preserve provides connectivity for wildlife from canyons leading from Chino Hills at the eastern Puente Hills west to the San Gabriel River and beyond. Each watershed was given a rating depending upon whether it had a high, medium, or low importance for wildlife connectivity. Table A-X shows the priority values associated with the different levels of importance.

**Table A-X: Wildlife Connectivity and Priority Values**


<b>Importance of Connectivity</b>	<b>Priority Value</b>
High Importance	10
Medium Importance	5
Low Importance	2

When all categories were designated, the priority values for each category were added together, resulting in a cumulative total to help create a basis for the restoration priorities. The resulting priority scores were then divided into five priority categories ranging from high to low. Table A-Y shows the priority categories and associated priority score totals. Figure A-8 shows the results of the weighted analysis for the overall priorities for restoration across the entire Preserve.

**Table A-Y: Restoration Priority Ranking Categories and Priority Score Ranges**


<b>Restoration Priority Ranking</b>	<b>Priority Score Ranges</b>
High Priority	70–94
Medium-High Priority	60–69
Medium Priority	50–59
Medium-Low Priority	40–49
Low Priority	0–39





Prepared By: L S A

FIGURE A-8



 PRESERVE BOUNDARY

RESTORATION PRIORITIES


 LOW RESTORATION PRIORITY

 MEDIUM-LOW RESTORATION PRIORITY

 MEDIUM RESTORATION PRIORITY

 MEDIUM-HIGH RESTORATION PRIORITY

 HIGH RESTORATION PRIORITY



Puente Hills Landfill  
Native Habitat Preservation Authority

Resource Management Plan

Overall Restoration Priorities

SOURCE: Aerial-EagleAerial (2003)  
I:\PUE430\GIS\Maps\Draft RMP\Appendices\FigA-8\_Overall\_Restunits\_Priorities.mxd (03/05/2007)



The weed polygons with priority ratings were then divided by restoration units. The restoration unit boundaries are based on watersheds. The restoration units were then ranked by calculating the percent of area occupied by weeds and multiplied by the categories in Table A-Z. The ranking calculation resulted in an ecological-based ranking, as shown in Figure A-9. However, some of the higher-ranked restoration units were not very feasible due to specific site conditions that were not reflected in the priority ranking system. The rankings of the restoration units were manually adjusted to account for this and could not be factored in by a calculation, as shown on Figure A-10. Specific electronic geographic information that contains all of these data will be provided separately to the Habitat Authority.


**Table A-Z: Restoration Unit Priority Ranking Multipliers by Percent of Weed Area**

<b>Percent of Restoration Unit Occupied by Weeds</b>	<b>Priority Ranking Multiplier</b>
0–20	1
20–40	1.1
40–60	1.3
60–80	1.4
80–100	1.5


## MANAGEMENT AREAS AND RESTORATION UNITS





As previously described, the Preserve has been divided into management areas based on ownership and adjacent communities. These management areas are discussed in the following section and restoration priorities have been calculated within each management area. Each of the management areas was analyzed and prioritized for restoration units by roughly-grouped watersheds to determine priority status for restoration. Named canyons and numbered watersheds are described in the following sections for each management area in order of the highest-priority restoration unit to the lowest priority. For each restoration unit, LSA developed a table identifying each weed polygon, the acreage, restoration priority rating, and proposed habitat to restore for polygons with a high to medium restoration priority. Where the weed polygon is one of the 93 soil sample areas, then LSA is confident of the determination of the habitat to be restored. Determination of the appropriate habitat included not only soils but also an analysis of remnant native species in the polygon, dominant weeds and cover, slope, aspect, and adjacent native habitats (specific electronic geographic information that contains all of these data will be provided separately to the Habitat Authority). If a weed polygon does not contain a specific associated soil sample, then the proposed habitat is followed by an asterisk (\*) indicating that it was determined based on general soil associations, rather than specific soil characteristics. For those specific invasive weed polygons extrapolated from BonTerra vegetation map, no habitat types were recommended. These polygons can be identified by the polygons in the 800 series. Additionally, analyses of remnant native species in the polygon, percent cover of dominant weeds, slope, aspect, and adjacent native habitats were used to suggest the appropriate habitat for restoration. It is LSA's recommendation that prior to restoration, soils be sampled in these polygons to confirm the appropriate habitat, as described previously. Because some weed polygons crossed watershed and management unit boundaries, some weed polygon numbers repeat within and across restoration units.





Prepared By: L S A



-  RESTORATION UNITS-LABELED WITH ECOLOGICAL PRIORITY RANKING (E.G., 13) AND NAME (E.G., W4)
-  HACIENDA HEIGHTS MANAGEMENT AREA
-  LA HABRA HEIGHTS MANAGEMENT AREA
-  WHITTIER MANAGEMENT AREA



 These parcels do not drain into the Arroyo Pescadero. However, for the purposes of the Habitat Restoration Plan, restoration data for these parcels is grouped with the Arroyo Pescadero Restoration Unit.

FIGURE A-9







LSA ASSOCIATES, INC.

PACIFIC CENTER  
703 PALOMAR AIRPORT RD., SUITE 260  
CARLSBAD, CALIFORNIA 92011760.931.5471 TEL  
760.918.2458  
FAXBERKELEY  
COLMA  
FORT COLLINSIRVINE  
PALM SPRINGS  
POINT RICHMONDRIVERSIDE  
ROCKLIN  
SAN LUIS OBISPO

August 4, 2009

Andrea Gullo  
Puente Hills Landfill Native Habitat Preservation Authority  
7702 Washington Avenue, Suite C  
Whittier, CA 90602

Subject: Summary of Focused Plant, Incidental and Protocol Survey Results (2008 and 2009)  
Puente Hills Landfill Native Habitat Preservation Authority Managed Lands  
City of Whittier Oil Exploration (LSA Project No. PUE0901)

Dear Ms. Gullo:

This letter report documents the results of incidental and protocol surveys for special-status plant and wildlife species conducted in 2008 and 2009 by LSA Associates, Inc. (LSA) for the proposed oil exploration activities on Puente Hills Landfill Native Habitat Preservation Authority (Habitat Authority) managed lands owned by and located in the City of Whittier, Los Angeles County, California (Figure 1; all figures are attached).

LSA conducted focused surveys for various special-status plants, protocol surveys for coastal California gnatcatcher (*Polioptila californica californica*) (CAGN) and least Bell's vireo (*Vireo bellii pusillus*) (LBVI) in accordance with United States Fish and Wildlife Service (USFWS) accepted survey protocols, and incidental surveys for cactus wren (*Campylorhynchus brunneicapillus*) and western spadefoot (*Spea hammondi*). Survey areas for 2008 and 2009 are shown in Figures 2 and 3.

The target special-status plant species, LBVI, and cactus wren were determined to be absent from the survey areas. Southern California black walnut (*Juglans californica*) was identified within the survey area (see Figure 2). No LBVI or CAGN were detected during protocol surveys in 2008, but in 2008, a Habitat Authority employee observed one CAGN in the 2008 survey area (see Figure 3).<sup>1</sup> In addition, one CAGN was observed during the final protocol survey conducted in 2009. Although western spadefoot was not observed during surveys, the species has the potential to occur within the survey areas.

## BACKGROUND

The 2008 survey area included five sites totaling approximately 113 acres under investigation for potential oil exploration activities within lands owned by the City of Whittier and managed by the Habitat Authority. In 2009, site boundaries were refined, two sites were eliminated, and the survey area was merged to include areas between the sites for a total survey area of approximately 209 acres. The 2008 and 2009 survey areas are located within Sections 22, 23, 25, and 26 of Township 2 South,

<sup>1</sup> David Moskovitz (USFWS-Permitted Individual to Survey for CAGN), personal communication, June 9 and 10, 2008.

Range 11 West, as shown on the United States Geological Survey (USGS) 7.5-minute *Whittier* and *La Habra*, California quadrangles (Figure 1).

The survey areas are characterized by portions of Arroyo Pescadero and La Cañada Verde drainages, adjacent hillsides, and access roads. Elevation ranges from approximately 300 to 1,000 feet above mean sea level. Vegetation communities within the survey areas primarily consist of coastal sage scrub, riparian scrub, nonnative grassland, ruderal vegetation, eucalyptus woodland (including some recently cleared areas), ornamental vegetation, and other previously disturbed areas. Soil associations mapped on the survey areas include Hanford Association (0 to 5 percent slopes), Perkins-Rincon Association (0 to 15 percent slopes), and Altamont-Diablo Association (30 to 60 percent slopes, eroded).<sup>1</sup>

Prior to conducting all surveys, LSA biologists reviewed appropriate literature and conducted a standard records search to determine whether special-status plant and wildlife species have been detected on or near the survey areas in the past. The literature review included various documents prepared by LSA for the Habitat Authority (including previous survey documents), as well as the California Natural Diversity Database (CNDDDB) maintained by the California Department of Fish and Game (CDFG), and the California Native Plant Society (CNPS) Online Inventory. The CNDDDB query included the *La Habra* and *Whittier*, California quadrangles, and the CNPS query included a 9-quad search of the surrounding quadrangles. LSA also reviewed the Resource Management Plan (RMP) prepared for the Habitat Authority in 2007, including the Sensitive Species Table in Appendix I, to further refine which special-status species might be present in the survey area.

Based on information gleaned in the above reviews, these species are known to occur within the elevation range and within vegetation communities of the survey areas: Plummer's mariposa lily (*Calochortus plummerae*), Robinson's peppergrass (*Lepidium virginicum* var. *robinsonii*), and Catalina mariposa lily (*Calochortus catalinae*), which are designated Special Plants by the CDFG and are CNPS Listed species, CAGN (federally threatened and California species of concern); LBVI (federally and California endangered); cactus wren (California species of concern); and western spadefoot (California species of concern). While the timing and methods of surveys focused on these species, all vertebrate wildlife and vascular plant species observed on the site were identified and recorded to document whether additional special-status species were present.

Prior to conducting protocol surveys, LSA sent 10-day notification letters on April 4, 2008, and on March 30, 2009, to the CDFG and the USFWS.

## METHODS

The 2008 survey area included five project sites totaling approximately 113 acres. See Table A-1 for a summary of each survey conducted in 2008.

---

<sup>1</sup> LSA, 2007. Resource Management Plan, Puente Hills Landfill Native Habitat Preservation Authority. Appendix B: Soil Taxonomy and Analysis.

**Table A-1: 2008 Survey Summary**

<b>Date</b>	<b>Surveyors</b>	<b>Plant</b>	<b>Protocol Wildlife</b>	<b>Incidental</b>
April 10, 2008	JH, DR	X		X
April 18, 2008	DR, JR	X		X
April 21, 2008	RE		X	X
May 1, 2008	RE		X	X
May 12, 2008	RE		X	X
May 22, 2008	RE		X	X
June 3, 2008	RE		X	X
June 4, 2008	JH, MW	X		X
June 5, 2008	JH, DR	X		X
June 13, 2008	RE		X	X
June 27, 2008	RE		X	X
July 9, 2008	RE		X	X

Surveyors: JH = Jim Harrison; DR = Dan Rosie; JR = Jodi Ross; RE = Richard Erickson;  
MW = Matthew Willis

The 2009 survey area included the consolidated survey area totaling approximately 209 acres. See Table A-2 for a summary of each survey conducted in 2009.

**Table A-2: 2009 Survey Summary**

<b>Date</b>	<b>Surveyors</b>	<b>Plant</b>	<b>Protocol Wildlife</b>	<b>Incidental</b>
April 2, 2009	DR, RS	X		X
April 3, 2009	DR, RS	X		X
April 15, 2009	MJB, RE		X	X
April 28, 2009	MJB, RE		X	X
May 8, 2009	MJB, RE		X	X
May 18, 2009	MJB, RE		X	X
May 28, 2009	RE		X	X
June 1, 2009	RE		X	X

Date	Surveyors	Plant	Protocol Wildlife	Incidental
June 5, 2009	SB, RS	X		X
June 8, 2009	MJB, RE		X	X
June 9, 2009	SB, RS	X		X
June 19, 2009	MJB, RE		X	X
June 29, 2009	MJB, RE		X	X

Surveyors: DR = Dan Rosie; RS = Robert Steers; MJB = Mark J. Billings; RE = Richard Erickson; SB = Sarah Barerra

### Focused Plant Surveys

LSA biologists Jim Harrison, Dan Rosie, and Robert Steers conducted focused plant surveys, with assistance from LSA biologists Jodi Ross, Matthew Willis, and Sarah Barerra.

The botanical surveys were conducted in accordance with the current CNPS Botanical Survey Guidelines dated June 2, 2001. Surveys were conducted by walking transects averaging approximately 50 feet wide, depending on visibility and habitat quality, throughout the survey area. The surveys were conducted during the expected flowering seasons for the target species (Plummer's mariposa lily, Robinson's peppergrass, and Catalina mariposa lily) in order to facilitate detection of these species even in a preflowering or postflowering state. In addition, steep slopes inaccessible by foot were surveyed with binoculars from the most practical vantage points.

A cumulative list of plant species identified during the 2008 and 2009 surveys appears in Table B (attached). Survey results are shown on Figure 2.

### Protocol Wildlife Surveys

All wildlife survey techniques followed available protocols established by the USFWS, other agencies, or interested parties. LSA biologists Richard Erickson and Mark J. Billings conducted protocol and incidental surveys pursuant to Federal Fish and Wildlife Permit TE777965-8 (April 8, 2008–April 7, 2012) and a temporary authorization letter from the CDFG (May 12, 2003–March 31, 2007; renewal request submitted March 26, 2007 and approved on May 4, 2007, extending coverage indefinitely) in lieu of a Memorandum of Understanding between LSA and CDFG.

Richard Erickson and Mark J. Billings conducted eight protocol LBVI and CAGN surveys from April 21 to July 9, 2008, and from April 28 to June 29, 2009. During each of the surveys, surveyors walked slowly along the edge of and, when appropriate, through riparian and coastal sage scrub habitat, listening for LBVI and CAGN. Surveyors played a tape recording of a CAGN periodically along the survey route during all of the surveys. With the aid of binoculars for viewing wildlife species, surveyors waited for several minutes after each playing to look and listen for both LBVI and CAGN. Survey results are shown on Figure 3.



## Incidental Wildlife Surveys

All LSA biologists conducted incidental surveys for cactus wren and western spadefoot concurrently with focused plant and wildlife surveys. They surveyed potentially suitable habitat for each species carefully to determine whether these species were present or absent.

A complete list of animals detected during the 2008 and 2009 surveys appears in Table C (attached).

## RESULTS

### Target Species

No threatened, endangered, or rare plant species, cactus wren, or western spadefoot were detected during surveys. No LBVI or CAGN were found during the 2008 surveys. However, a single CAGN was observed in the 2008 survey area by Habitat Authority employee David Moskovitz on June 9 and 10, 2008. A search of the same area by Mr. Moskovitz and Mr. Erickson on June 13, 2008, was unsuccessful. The bird was most likely a recently fledged juvenile dispersing from its natal territory. No LBVI were found during the 2009 surveys. In 2009, a single young male CAGN was seen in the 2009 survey area by Mr. Erickson on June 29. This bird was most likely a recently fledged juvenile dispersing from its natal territory. Figure 3 illustrates the location of these individuals and of potentially suitable nesting habitat.

### Other Species

**Plants.** A small stand (five individuals) of Southern California black walnut (*Juglans californica*) was found in the drainage that runs parallel and east of Catalina Avenue, about 0.15 mile north of the entrance gate. This small stand is a component of the riparian vegetation that occurs in the drainage. It does not constitute a California walnut woodland or walnut forest, which are vegetation types of interest to the CDFG. Southern California black walnut is on the CNPS 4.2 list. CNPS List 4 is only a “watch list” and species on this list do not appear on CNPS or CNDDDB searches by USGS quads. This species has no State or federal status and is generally not afforded the same level of protection as species that are listed as threatened or endangered, but it is included in the Resource Management Plan for the preserve. Figure 2 illustrates the location of this species within the survey area.

**Wildlife.** The following special-status wildlife species, which are considered to be California Species of Concern (CSC) or California “Special Animals” (CSA) according to the CDFG, were observed. Note that certain species are only considered special-status when nesting; although nesting was not necessarily observed during surveys, it is possible that these species may nest in or near the survey area.

- Western whiptail (*Aspidoscelis tigris*) (CSA)
- Cooper’s hawk (*Accipiter cooperii*) (CSA when nesting)
- Sharp-shinned hawk (*Accipiter striatus*) (CSA when nesting)
- Vaux’s swift (*Chaetura vauxi*) (CSC when nesting)

- Costa's hummingbird (*Calypte costae*) (CSA when nesting)
- California horned lark (*Eremophila alpestris actia*)<sup>1</sup> (CSA)
- Rufous/Allen's hummingbird (*Selasphorus rufus/sasin*) (both CSA when nesting)
- Nuttall's woodpecker (*Picoides nuttallii*) (CSA when nesting)
- Yellow warbler (*Dendroica petechia*) (CSC when nesting)
- Yellow-breasted chat (*Icteria virens*) (CSC when nesting)
- Southern California rufous-crowned sparrow (*Aimophila ruficeps canescens*) (CSA when nesting)
- Brewer's sparrow (*Spizella breweri*) (CSA when nesting)
- Chipping sparrow (*Spizella passerina*) (CSA when nesting)
- San Diego desert woodrat (*Neotoma lepida intermedia*) (CSC)

The following species are considered to be CSA only when rookery sites are present. These species were observed during surveys, but no rookery sites were detected:

- Double-crested cormorant (*Phalacrocorax auritus*)
- Great blue heron (*Ardea herodias*)
- Great egret (*Ardea alba*)
- Caspian tern (*Hydroprogne caspia*)

## CONCLUSIONS

Based on the results of the focused plant surveys, LSA concludes that special-status plant species are not present within the survey areas.

The CAGN Mr. Moskovitz observed in 2008 and the CAGN observed during protocol surveys in 2009 appeared to be passing through the area and not to be resident birds. At this time, CAGN are considered not to be nesting within the survey areas although potentially suitable nesting habitat is present within and adjacent to the survey areas.

Cactus wren was not observed during surveys and is considered unlikely to occur, given the marginal suitability of the habitat within the survey areas.

LBVI was not observed during protocol surveys. This species is considered absent from the survey areas at this time although potentially suitable nesting habitat occurs.

---

<sup>1</sup> Both *Eremophila alpestris actia* and *E. a. ammophila* are known from the region. The bird observed during surveys may have been either subspecies; however, *E. a. actia* has been recorded on the site previously as documented in Appendix I of the RMP.

Western spadefoot was not observed during the surveys; however, suitable habitat occurs throughout the survey areas, and this species is known to occur within the vicinity of the survey areas. Therefore, this species is considered potentially present within the survey areas.

If you have any questions or require additional information, please feel free to call me at (760) 931-5471 or e-mail me at [Adrienne.Bezley@lsa-assoc.com](mailto:Adrienne.Bezley@lsa-assoc.com).

Sincerely,

**LSA ASSOCIATES, INC.**



Adrienne Bezley  
Senior Biologist

Attachments: A: Figures and Tables  
    Figure 1: Regional Location  
    Figure 2: 2008 and 2009 Plant Survey Areas and Results  
    Figure 3: 2008 and 2009 Wildlife Survey Areas and Results  
    Table B: Plant Species Observed  
    Table C: Wildlife Species Detected  
B: Focused Plant Survey Report  
C: 2008 and 2009 Protocol Least Bell's Vireo and Coastal California Gnatcatcher Survey Reports

## REFERENCES

- California Department of Fish and Game, Natural Heritage Division, Natural Diversity Database. 2008. RareFind Version 3.1.0. Records search executed April 8, 2008, covering the USGS 7.5-minute series topographic map, La Habra and Whittier, California quadrangles. Sacramento, California: The Resources Agency. Commercial version dated February 2, 2008.
- California Native Plant Society. 2001. CNPS Botanical Survey Guidelines. Revised June 2, 2001.
- California Native Plant Society (CNPS). 2008. Inventory of Rare and Endangered Plants (online edition, v7-08b). California Native Plant Society. Sacramento, CA. Accessed on April 8, 2008 from <http://www.cnps.org/inventory>.
- Hickman, J.C. (ed.). 1996. The Jepson Manual: Higher Plants of California. University of California Press, Berkeley.
- Jepson Online Interchange for California Floristics. 2009. Accessed on June 23, 2009. [<http://ucjeps.berkeley.edu/interchange.html>]
- Ljubenkov, J.A.S., and T.S. Ross. 2002. An Annotated Checklist of the Vascular Plants of the Whittier Hills, Los Angeles County, California. *Crossoma* 27(1), Spring-Summer 2001 (issued August 2002).
- LSA Associates, Inc. 2006. Botanical Survey Report 2005. Prepared for the Puente Hills Landfill Native Habitat Preservation Authority. January 13 2006.
- LSA Associates, Inc. 2007. Resource Management Plan. Prepared for the Puente Hills Landfill Native Habitat Preservation Authority. July 2007.



## ATTACHMENT A: FIGURES AND TABLES







FIGURE 1

LSA

LEGEND


- 2009 Survey Area (approx. 209 acres)
- 2008 Survey Area (approx. 111 acres)



Puente Hills Landfill Native Habitat Preservation Authority  
 City of Whittier Oil Exploration  
 Project Location

SOURCE: USGS 7.5' QUAD - WHITTIER (81) AND LA HABRA (81); CALIF.







LSA

LEGEND

- 2009 Survey Areas (approx. 209 acres) Southern California Black Walnut (*Juglans californica*)
- 2008 Survey Areas (approx. 111 acres)
- Trees mapped within Survey Area
- Trees observed outside Survey Area (approx. 10c)




SOURCE: DigitalGlobe (April 2008)

I:\PUE0901\GIS\Reports\Survey\_Summary\Fig2\_Plant\_Survey\_Results.mxd (08/05/2009)



FIGURE 2

Puente Hills Landfill Native Habitat Preservation Authority  
City of Whittier Oil Exploration  
2008 and 2009 Plant Survey Areas and Results





LSA



SOURCE: DigitalGlobe (April 2008)

I:\PUE0901\GIS\Reports\Bird\_Survey\Fig3\_Bird\_Survey\_Results.mxd (08/04/2009)







-  Coastal California Gnatcatcher Observed June 29, 2009
-  Coastal California Gnatcatcher Observed June 9 & 10, 2008
-  Most Suitable Coastal California Gnatcatcher Habitat
-  Potential Least Bell's Vireo Habitat
-  2009 Survey Area
-  2008 Survey Area

FIGURE 3

Puente Hills Landfill Native Habitat Preservation Authority  
 City of Whittier Oil Exploration  
 2008 and 2009 Wildlife Survey Areas and Results



## TABLE B

### VASCULAR PLANT SPECIES OBSERVED

The following vascular plant species were observed in the survey areas by various biologists during the course of on-site surveys in 2008 and 2009. Family- and species-level nomenclature follows the Jepson Manual. For species nomenclature only, current name changes on the Jepson Interchange are also used.

\*Introduced nonnative species

\*\*Special-status species

Scientific Name	Common Name
<b>ANGIOSPERMAE: DICOTYLEDONAE</b>	<b>DICOT FLOWERING PLANTS</b>
<b>Anacardiaceae</b>	<b>Sumac Family</b>
<i>Malosma laurina</i>	Laurel sumac
<i>Rhus integrifolia</i>	Lemonade berry
<i>Rhus ovata</i>	Sugar bush
* <i>Schinus molle</i>	Peruvian pepper tree
* <i>Schinus terebinthifolius</i>	Brazilian pepper tree
<i>Toxicodendron diversilobum</i>	Poison oak
<b>Apiaceae</b>	<b>Carrot Family</b>
* <i>Conium maculatum</i>	Poison hemlock
<i>Daucus pusillus</i>	American wild carrot
* <i>Foeniculum vulgare</i>	Sweet fennel
<b>Apocynaceae</b>	<b>Dogbane Family</b>
* <i>Vinca major</i>	Periwinkle
<b>Asclepiadaceae</b>	<b>Milweed Family</b>
<i>Asclepias californica</i>	California milkweed
<b>Asteraceae</b>	<b>Sunflower Family</b>
<i>Ambrosia acanthicarpa</i>	Annual sandbar
<i>Ambrosia psilostachya</i>	Western ragweed
<i>Artemisia californica</i>	California sagebrush
<i>Artemisia douglasiana</i>	California mugwort
<i>Baccharis emoryi</i>	Emory's baccharis
<i>Baccharis pilularis</i>	Coyote bush
<i>Baccharis salicifolia</i>	Mule fat
* <i>Carduus pycnocephalus</i>	Italian thistle
* <i>Centaurea melitensis</i>	Tocalote
* <i>Cirsium vulgare</i>	Bull thistle
<i>Corethrogyne filaginifolia</i>	California aster
<i>Deinandra fasciculata</i>	Fascicled tarweed
<i>Encelia californica</i>	California bush sunflower
<i>Gutierrezia californica</i>	California matchweed
* <i>Hedypnois cretica</i>	Crete hedypnois
<i>Helianthus annuus</i>	Annual sunflower
<i>Heterotheca grandiflora</i>	Telegraph weed

Scientific Name	Common Name
<i>Isocoma menziesii</i>	Coastal goldenbush
* <i>Lactuca serriola</i>	Prickly lettuce
<i>Corethrogyne filaginifolia</i>	Common sand aster
<i>Logfia filaginoides</i>	California filago
<i>Malacothrix saxatilis</i> var. <i>tenuifolia</i>	Cliff malacothrix
* <i>Picris echioides</i>	Bristly ox tongue
<i>Pseudognaphalium biolettii</i>	Two-color rabbit-tobacco
<i>Pseudognaphalium californicum</i>	California rabbit-tobacco
* <i>Pseudognaphalium luteoalbum</i>	Jersey cudweed
<i>Pseudognaphalium microcephalum</i>	San Diego rabbit-tobacco
* <i>Senecio vulgaris</i>	Common groundsel
* <i>Silybum marianum</i>	Milk thistle
* <i>Sonchus asper</i>	Common sow thistle
* <i>Sonchus oleraceus</i>	Common sow thistle
<i>Stephanomeria virgata</i>	Tall wreath plant
<i>Xanthium strumarium</i>	Cocklebur
<b>Brassicaceae</b>	<b>Mustard Family</b>
* <i>Brassica nigra</i>	Black mustard
* <i>Hirschfeldia incana</i>	Summer mustard
* <i>Raphanus sativus</i>	Wild radish
* <i>Sisymbrium</i> sp.	Mustard
* <i>Sisymbrium irio</i>	London rocket
<b>Cactaceae</b>	<b>Cactus Family</b>
<i>Opuntia littoralis</i>	Coastal prickly pear
<b>Caprifoliaceae</b>	<b>Honeysuckle Family</b>
<i>Sambucus mexicana</i>	Mexican elderberry
<b>Caryophyllaceae</b>	<b>Pink Family</b>
<i>Stellaria media</i>	Chickweed
<b>Chenopodiaceae</b>	<b>Goosefoot Family</b>
* <i>Atriplex semibaccata</i>	Australian saltbush
* <i>Chenopodium album</i>	Lamb's quarters
<i>Chenopodium berlandieri</i>	Pitseed goosefoot
* <i>Chenopodium murale</i>	Nettle leaf goosefoot
* <i>Salsola tragus</i>	Russian thistle
<b>Convolvulaceae</b>	<b>Morning-Glory Family</b>
<i>Calystegia macrostegia</i>	Morning-glory
<b>Crassulaceae</b>	<b>Stonecrop Family</b>
<i>Crassula connata</i>	Sand pygmy-stonecrop
<b>Cucurbitaceae</b>	<b>Gourd Family</b>
<i>Cucurbita foetidissima</i>	Coyote melon
<i>Marah macrocarpus</i>	Man root
<b>Euphorbiaceae</b>	<b>Spurge Family</b>
<i>Chamaesyce albomarginata</i>	Rattlesnake weed
<i>Chamaesyce</i> sp.	Spurge
* <i>Ricinis communis</i>	Castor bean
<b>Fabaceae</b>	<b>Legume Family</b>
* <i>Acacia cyclops</i>	Coastal wattle
* <i>Acacia longifolia</i>	Golden wattle
<i>Lotus salsuginosus</i> var. <i>salsuginosus</i>	Coastal lotus

Scientific Name	Common Name
<i>Lotus scoparius</i>	Deer weed
<i>Lupinus microcarpus</i> var. <i>densiflorus</i>	Dense-flowered chick lupine
<i>Lupinus succulentus</i>	Arroyo lupine
* <i>Medicago polymorpha</i>	Bur clover
* <i>Melilotus indica</i>	Yellow sweet clover
<b>Fagaceae</b>	<b>Oak Family</b>
<i>Quercus agrifolia</i>	Coast live oak
<b>Geraniaceae</b>	<b>Geranium Family</b>
* <i>Erodium cicutarium</i>	Red-stemmed filaree
* <i>Erodium moschatum</i>	White-stemmed filaree
<b>Grossulariaceae</b>	<b>Gooseberry Family</b>
<i>Ribes speciosum</i>	Fuchsia flowered gooseberry
<b>Hydrophyllaceae</b>	<b>Waterleaf Family</b>
<i>Emmenanthe penduliflora</i>	Whispering bells
<i>Eucrypta chrysanthemifolia</i>	Common eucrypta
<i>Phacelia cicutaria</i> var. <i>hispida</i>	Caterpillar phacelia
<i>Phacelia minor</i>	California bluebell
<i>Phacelia parryi</i>	Parry's phacelia
<i>Phacelia ramosissima</i>	Branching phacelia
<i>Phacelia tanacetifolia</i>	Tansy leafed phacelia
<i>Pholistoma auritum</i> var. <i>auritum</i>	Blue fiesta flower
<b>Juglandaceae</b>	<b>Walnut Family</b>
** <i>Juglans californica</i>	Southern California walnut
<b>Lamiaceae</b>	<b>Mint Family</b>
* <i>Marrubium vulgare</i>	Horehound
<i>Salvia apiana</i>	White sage
<i>Salvia leucophylla</i> X <i>apiana</i>	White/purple sage hybrid
<i>Salvia leucophylla</i>	Purple sage
<i>Salvia mellifera</i>	Black sage
<b>Malvaceae</b>	<b>Mallow Family</b>
<i>Malacothamnus fasciculatus</i>	Lax-flowered mallow
* <i>Malva parviflora</i>	Cheeseweed
* <i>Malva sylvestris</i>	High mallow
<b>Myrtaceae</b>	<b>Myrtle Family</b>
* <i>Eucalyptus</i> sp.	Gum
<b>Nyctaginaceae</b>	<b>Four-O'Clock Family</b>
<i>Mirabilis laevis</i>	Wishbone bush
<b>Onagraceae</b>	<b>Evening Primrose Family</b>
<i>Camissonia californica</i>	California suncup
<i>Clarkia bottae</i>	Botta's clarkia
<b>Oxalidaceae</b>	<b>Oxalis Family</b>
* <i>Oxalis pes-caprae</i>	Bermuda buttercup
<b>Platanaceae</b>	<b>Sycamore Family</b>
<i>Platanus racemosa</i>	California sycamore
<b>Polygonaceae</b>	<b>Buckwheat Family</b>
<i>Eriogonum fasciculatum</i>	California buckwheat
* <i>Rumex crispus</i>	Curly dock
<b>Portulacaceae</b>	<b>Purselane Family</b>
<i>Claytonia perfoliata</i>	Miner's lettuce

Scientific Name	Common Name
<b>Primulaceae</b>	<b>Primrose Family</b>
* <i>Anagallis arvensis</i>	Scarlet pimpernel
<b>Rosaceae</b>	<b>Rose Family</b>
<i>Heteromeles arbutifolia</i>	Toyon
<i>Prunus ilicifolia</i>	Hollyleaf cherry
<b>Rubiaceae</b>	<b>Madder Family</b>
<i>Galium angustifolium</i>	Chaparral bedstraw
<i>Galium aparine</i>	Common bedstraw
<b>Salicaceae</b>	<b>Willow Family</b>
<i>Salix exigua</i>	Narrowleaf willow
<i>Salix gooddingii</i>	Black willow
<i>Salix laevigata</i>	Red willow
<i>Salix lasiolepis</i>	Arroyo willow
<b>Scrophulariaceae</b>	<b>Figwort Family</b>
<i>Keckiella cordifolia</i>	Heart-leaved penstemon
<i>Mimulus aurantiacus</i>	Orange bush monkey flower
<i>Scrophularia californica</i>	California figwort
* <i>Verbascum virgatum</i>	Wandmullein
<b>Solanaceae</b>	<b>Nightshade Family</b>
* <i>Nicotiana glauca</i>	Tree tobacco
* <i>Solanum americanum</i>	White nightshade
<i>Solanum douglasii</i>	Douglas' nightshade
<b>Tamaricaceae</b>	<b>Tamarisk Family</b>
* <i>Tamarix ramosissima</i>	Mediterranean tamarisk
<b>Tropaeolaceae</b>	<b>Nightshade Family</b>
* <i>Tropaeolum majus</i>	Garden nasturtium
<b>Ulmaceae</b>	<b>Elm Family</b>
* <i>Ulmus parvifolia</i>	Chinese elm
<b>Urticaceae</b>	<b>Nettle Family</b>
<i>Urtica dioica</i> ssp. <i>holosericea</i>	Hoary nettle
* <i>Urtica urens</i>	Dwarf nettle
<i>Parietaria hespera</i>	Western pellitory
<b>Verbenaceae</b>	<b>Vervain Family</b>
<i>Verbena lasiostachys</i>	Western verbena
<b>ANGIOSPERMAE: MONOCOTYLEDONAE</b>	<b>MONOCOT FLOWERING PLANTS</b>
<b>Arecaceae</b>	<b>Palm Family</b>
* <i>Washingtonia robusta</i>	Mexican fan palm
<b>Iridaceae</b>	<b>Iris Family</b>
<i>Sisyrinchium bellum</i>	Blue eyed grass
<b>Liliaceae</b>	<b>Lily Family</b>
<i>Dichelostemma capitatum</i>	Blue dicks
<b>Poaceae</b>	<b>Grass Family</b>
* <i>Avena barbata</i>	Slender wild oat
* <i>Avena fatua</i>	Wild oat
* <i>Bromus diandrus</i>	Ripgut brome
* <i>Bromus hordeaceus</i>	Soft chess
* <i>Bromus madritensis</i> ssp. <i>rubens</i>	Red brome
* <i>Cortaderia jubata</i>	Pampas grass
* <i>Cynodon dactylon</i>	Bermuda grass



<b>Scientific Name</b>	<b>Common Name</b>
* <i>Hordeum murinum</i> ssp. <i>leporinum</i>	Foxtail barley
<i>Leymus condensatus</i>	Giant wildrye
<i>Leymus triticoides</i>	Creeping wild rye
* <i>Lolium multiflorum</i>	Italian ryegrass
<i>Nassella pulchra</i>	Purple needlegrass
<i>Nassella lepida</i>	Foothill needlegrass
* <i>Pennisetum setaceum</i>	African fountain grass
* <i>Piptatherum miliaceum</i>	Smilo grass
* <i>Schismus barbatus</i>	Schismus
* <i>Vulpia myuros</i>	Foxtail fescue

## TABLE C

### WILDLIFE SPECIES DETECTED

This is a list of the conspicuous aerial insects, amphibians, reptiles, birds, and mammals noted in the survey areas by LSA biologists. Presence may be noted if a species is seen or heard, or identified by the presence of tracks, scat, or other signs.

- \* Species not native to the area
- \*\* Special-status species (note that some species are only considered special-status if nesting colonies are present)

Scientific Name	Common Name
<b>ANISOPTERA</b>	<b>TYPICAL DRAGONFLIES</b>
<b>Libellulidae</b>	<b>Cruisers, Emeralds, Baskettails, and Skimmers</b>
<i>Libellula saturata</i>	Flame skimmer
<i>Pantala hymenaea</i>	Spot-winged glider
<i>Pantala hymenaea</i>	Spot-winged glider
<b>LEPIDOPTERA</b>	<b>BUTTERFLIES</b>
<b>Papilionidae</b>	<b>Swallowtails</b>
<i>Papilio rutulus</i>	Western tiger swallowtail
<b>Pieridae</b>	<b>Whites and Sulphurs</b>
<i>Pontia protodice</i>	Checkered white
* <i>Pieris rapae</i>	Cabbage white
<i>Anthocharis sara</i>	Sara orangetip
<i>Colias eurytheme</i>	Orange sulphur
<b>Lycaenidae</b>	<b>Gossamer-Wing Butterflies</b>
<i>Leptotes marina</i>	Marine blue
<i>Plebejus acmon</i>	Acmon blue
<b>Nymphalidae</b>	<b>Brush-Footed Butterflies</b>
<i>Agraulis vanillae</i>	Gulf fritillary
<i>Nymphalis antiopa</i>	Mourning cloak
<i>Vanessa cardui</i>	Painted lady
<i>Vanessa annabella</i>	West coast lady
<i>Vanessa atalanta</i>	Red admiral
<b>AMPHIBIA</b>	<b>AMPHIBIANS</b>
<b>Hylidae</b>	<b>Treefrogs and Relatives</b>
<i>Pseudacris hypochondriaca</i>	Baja California treefrog
<b>REPTILIA</b>	<b>REPTILES</b>
<b>Phrynosomatidae</b>	<b>Phrynosomatid Lizards</b>
<i>Sceloporus occidentalis</i>	Western fence lizard
<b>Teiidae</b>	<b>Whiptails and Relatives</b>
** <i>Aspidoscelis tigris</i>	Western whiptail

Scientific Name	Common Name
<b>Colubridae</b>	<b>Colubrid Snakes</b>
<i>Tantilla planiceps</i>	California black-headed snake
<b>Viperidae</b>	<b>Vipers</b>
<i>Crotalus oreganus</i>	Western rattlesnake
<b>AVES</b>	<b>BIRDS</b>
<b>Anatidae</b>	<b>Ducks, Geese, and Swans</b>
<i>Anas platyrhynchos</i>	Mallard
<b>Odontophoridae</b>	<b>New World Quail</b>
<i>Callipepla californica</i>	California quail
<b>Phalacrocoracidae</b>	<b>Cormorants</b>
** <i>Phalacrocorax auritus</i>	Double-crested cormorant
<b>Ardeidae</b>	<b>Hérons, Bitterns, and Allies</b>
** <i>Ardea herodias</i>	Great blue heron
** <i>Ardea alba</i>	Great egret
<b>Cathartidae</b>	<b>New World Vultures</b>
<i>Cathartes aura</i>	Turkey vulture
<b>Accipitridae</b>	<b>Hawks, Kites, Eagles, and Allies</b>
** <i>Accipiter striatus</i>	Sharp-shinned hawk
** <i>Accipiter cooperii</i>	Cooper's hawk
<i>Buteo lineatus</i>	Red-shouldered hawk
<i>Buteo jamaicensis</i>	Red-tailed hawk
<b>Falconidae</b>	<b>Caracaras and Falcons</b>
<i>Falco sparverius</i>	American kestrel
<b>Laridae</b>	<b>Gulls, Terns, and Skimmers</b>
<i>Larus californicus</i>	California gull
<i>Larus occidentalis</i>	Western gull
<i>Larus glaucescens</i>	Glaucous-winged gull
** <i>Hydroprogne caspia</i>	Caspian tern
<b>Columbidae</b>	<b>Pigeons and Doves</b>
* <i>Columba livia</i>	Rock (Feral) pigeon
<i>Patagioenas fasciata</i>	Band-tailed pigeon
<i>Zenaida macroura</i>	Mourning dove
<b>Psittacidae</b>	<b>Lories, Parakeets, Macaws, and Parrots</b>
* <i>Brotogeris chiriri</i>	Yellow-chevroned parakeet
* <i>Amazona viridigenalis</i>	Red-crowned parrot
<b>Strigidae</b>	<b>Typical Owls</b>
<i>Bubo virginianus</i>	Great horned owl
<b>Apodidae</b>	<b>Swifts</b>
** <i>Chaetura vauxi</i>	Vaux's swift
<i>Aeronautes saxatilis</i>	White-throated swift
<b>Trochilidae</b>	<b>Hummingbirds</b>
<i>Archilochus alexandri</i>	Black-chinned hummingbird
<i>Calypte anna</i>	Anna's hummingbird
** <i>Calypte costae</i>	Costa's hummingbird
** <i>Selasphorus rufus/sasin</i>	Rufous/Allen's hummingbird
<b>Picidae</b>	<b>Woodpeckers and Allies</b>
<i>Melanerpes formicivorus</i>	Acorn woodpecker
<i>Sphyrapicus</i> sp.	Sapsucker sp.



Scientific Name	Common Name
** <i>Picoides nuttallii</i>	Nuttall's woodpecker
<i>Picoides pubescens</i>	Downy woodpecker
<i>Colaptes auratus</i>	Northern flicker
<b>Tyrannidae</b>	<b>Tyrant Flycatchers</b>
<i>Contopus sordidulus</i>	Western wood-pewee
<i>Empidonax difficilis</i>	Pacific-slope flycatcher
<i>Sayornis nigricans</i>	Black phoebe
<i>Myiarchus cinerascens</i>	Ash-throated flycatcher
<i>Tyrannus vociferans</i>	Cassin's kingbird
<i>Tyrannus verticalis</i>	Western kingbird
<b>Vireonidae</b>	<b>Vireos</b>
<i>Vireo huttoni</i>	Hutton's vireo
<i>Vireo gilvus</i>	Warbling vireo
<b>Corvidae</b>	<b>Crows and Jays</b>
<i>Aphelocoma californica</i>	Western scrub-jay
<i>Corvus brachyrhynchos</i>	American crow
<i>Corvus corax</i>	Common raven
<b>Alaudidae</b>	<b>Larks</b>
** <i>Eremophila alpestris</i>	Horned lark
<b>Hirundinidae</b>	<b>Swallows</b>
<i>Stelgidopteryx serripennis</i>	Northern rough-winged swallow
<i>Petrochelidon pyrrhonota</i>	Cliff swallow
<i>Hirundo rustica</i>	Barn swallow
<b>Aegithalidae</b>	<b>Long-Tailed Tits and Bushtits</b>
<i>Psaltriparus minimus</i>	Bushtit
<b>Troglodytidae</b>	<b>Wrens</b>
<i>Thryomanes bewickii</i>	Bewick's wren
<i>Troglodytes aedon</i>	House wren
<b>Turdidae</b>	<b>Thrushes</b>
<i>Sialia mexicana</i>	Western bluebird
<i>Catharus ustulatus</i>	Swainson's thrush
<i>Catharus guttatus</i>	Hermit thrush
<i>Turdus migratorius</i>	American robin
<b>Timaliidae</b>	<b>Babblers</b>
<i>Chamaea fasciata</i>	Wrentit
<b>Mimidae</b>	<b>Mockingbirds and Thrashers</b>
<i>Mimus polyglottos</i>	Northern mockingbird
<i>Toxostoma redivivum</i>	California thrasher
<b>Bombycillidae</b>	<b>Waxwings</b>
<i>Bombycilla cedrorum</i>	Cedar waxwing
<b>Ptilonotidae</b>	<b>Silky-flycatchers</b>
<i>Phainopepla nitens</i>	Phainopepla
<b>Parulidae</b>	<b>Wood Warblers</b>
<i>Vermivora celata</i>	Orange-crowned warbler
<i>Vermivora ruficapilla</i>	Nashville warbler
** <i>Dendroica petechia</i>	Yellow warbler
<i>Dendroica coronata</i>	Yellow-rumped warbler
<i>Dendroica nigrescens</i>	Black-throated gray warbler
<i>Dendroica townsendi</i>	Townsend's warbler

Scientific Name	Common Name
<i>Oporornis tolmiei</i>	MacGillivray's warbler
<i>Dendroica occidentalis</i>	Hermit warbler
<i>Geothlypis trichas</i>	Common yellowthroat
<i>Wilsonia pusilla</i>	Wilson's warbler
** <i>Icteria virens</i>	Yellow-breasted chat
<b>Thraupidae</b>	<b>Tanagers</b>
<i>Piranga ludoviciana</i>	Western tanager
<b>Emberizidae</b>	<b>Emberizids</b>
<i>Pipilo maculatus</i>	Spotted towhee
<i>Pipilo crissalis</i>	California towhee
** <i>Aimophila ruficeps canescens</i>	Southern California rufous-crowned sparrow
** <i>Spizella breweri</i>	Brewer's sparrow
** <i>Spizella passerina</i>	Chipping sparrow
<i>Passerella iliaca</i>	Fox sparrow
<i>Melospiza melodia</i>	Song sparrow
<i>Zonotrichia leucophrys</i>	White-crowned sparrow
<i>Zonotrichia atricapilla</i>	Golden-crowned sparrow
<b>Cardinalidae</b>	<b>Cardinals, Saltators, and Allies</b>
<i>Pheucticus ludovicianus</i>	Rose-breasted grosbeak
<i>Pheucticus melanocephalus</i>	Black-headed grosbeak
<i>Passerina caerulea</i>	Blue grosbeak
<i>Passerina amoena</i>	Lazuli bunting
<b>Icteridae</b>	<b>Blackbirds</b>
<i>Euphagus cyanocephalus</i>	Brewer's blackbird
<i>Agelaius phoeniceus</i>	Red-winged blackbird
<i>Molothrus ater</i>	Brown-headed cowbird
<i>Icterus cucullatus</i>	Hooded oriole
<b>Icteridae (cont'd)</b>	<b>Blackbirds</b>
<i>Icterus bullockii</i>	Bullock's oriole
<b>Fringillidae</b>	<b>Fringilline and Cardueline Finches and Allies</b>
<i>Carpodacus mexicanus</i>	House finch
<i>Carduelis psaltria</i>	Lesser goldfinch
<i>Carduelis tristis</i>	American goldfinch
<b>Passeridae</b>	<b>Old World Sparrows</b>
* <i>Passer domesticus</i>	House sparrow
<b>Estrildidae</b>	<b>Estrildid Finches</b>
* <i>Lonchura punctulata</i>	Nutmeg mannikin
<b>MAMMALIA</b>	<b>MAMMALS</b>
<b>Sciuridae</b>	<b>Squirrels, Chipmunks, and Marmots</b>
* <i>Sciurus niger</i>	Eastern fox squirrel
<i>Spermophilus beecheyi</i>	California ground squirrel
<b>Geomyidae</b>	<b>Pocket Gophers</b>
<i>Thomomys bottae</i>	Botta's pocket gopher
<b>Cricetidae</b>	<b>Hamsters, Voles, Lemmings, and New World Rats and Mice</b>
** <i>Neotoma lepida intermedia</i>	San Diego desert woodrat
<i>Neotoma macrotis</i>	Big-eared woodrat
<b>Leporidae</b>	<b>Rabbits and Hares</b>

Scientific Name	Common Name
<i>Sylvilagus audubonii</i>	Audubon's cottontail
<b>Felidae</b>	<b>Cats</b>
<i>Lynx rufus</i>	Bobcat
<b>Canidae</b>	<b>Foxes, Wolves, and Allies</b>
<i>Canis latrans</i>	Coyote
<b>Cervidae</b>	<b>Deer, Elk, and Allies</b>
<i>Odocoileus hemionus</i>	Mule deer

**Taxonomy and nomenclature are based on the following:**

Damselflies and dragonflies: Manolis, T. (2003, Dragonflies and Damselflies of California, University of California Press, Berkeley).

Butterflies: North American Butterfly Association (2001, NABA checklist and English Names of North American Butterflies, Second Edition, North American Butterfly Association, Morristown, New Jersey).

Amphibians and reptiles: Crother, B.I. ed. (2008. Scientific and Standard English Names of Amphibians and Reptiles of North America North of Mexico. Herpetological Circular 37) for species taxonomy and nomenclature; Stebbins, R.C. (2003, A Field Guide to Western Reptiles and Amphibians, third edition, Houghton Mifflin, Boston) for sequence and higher order taxonomy.

Birds: American Ornithologists' Union (1998, The A.O.U. Checklist of North American Birds, Seventh Edition, American Ornithologists' Union, Washington D.C.; and 2000, 2002, 2003, 2004, 2005, 2006, and 2007 supplements; see <http://aou.org.whsites.net/checklist/index.php3>).

Mammals: Wilson, D.E., and D.M. Reeder, eds. (2005. Mammal Species of the World, 3rd ed. Johns Hopkins University Press, Baltimore, Maryland; see <http://nmmhgoph.si.edu/msw/>).



## **ATTACHMENT B: FOCUSED PLANT SURVEY REPORT**

**ATTACHMENT C: 2008 AND 2009 PROTOCOL LEAST BELL'S  
VIREO AND COASTAL CALIFORNIA GNATCATCHER SURVEY  
REPORTS**



LSA ASSOCIATES, INC.

PACIFIC CENTER  
703 PALOMAR AIRPORT RD., SUITE 260  
CARLSBAD, CALIFORNIA 92011

760.931.5471 TEL  
760.918.2458  
FAX

BERKELEY  
COLMA  
FORT COLLINS

IRVINE  
PALM SPRINGS  
POINT RICHMOND

RIVERSIDE  
ROCKLIN  
SAN LUIS OBISPO

August 4, 2009

Andrea Gullo  
Puente Hills Landfill Native Habitat Preservation Authority  
7702 Washington Avenue, Suite C  
Whittier, CA 90602

Subject: Focused Survey Results  
Special-Status Plant Species  
City of Whittier Oil Exploration (LSA Project No. PUE0901)

Dear Ms. Gullo:

This letter report documents the results of focused plant surveys conducted by LSA Associates, Inc. (LSA) for the proposed oil exploration activities within lands managed by the Puente Hills Landfill Native Habitat Preservation Authority (Habitat Authority), owned by and located in the City of Whittier, Los Angeles County, California (Figure 1; all figures are attached).

After two years of surveys (2008 and 2009), special-status plant species were determined to be absent from the survey areas.

## BACKGROUND

In 2008, the survey area included five sites totaling approximately 113 acres under investigation for potential oil exploration activities. In 2009, the survey area was larger, totaling approximately 209 acres that overlapped much of the land surveyed in 2008 (Figure 2). Both the 2008 and 2009 survey areas are located within Sections 22, 23, 25, and 26 of Township 2 South, Range 11 West, as shown on the United States Geological Survey (USGS) 7.5-minute *Whittier* and *La Habra, California* quadrangles.

The survey areas are characterized by portions of Arroyo Pescadero and La Cañada Verde drainages, adjacent hillsides, and access roads. Elevations range from approximately 300 to 1,000 feet above sea level. Vegetation types within the survey areas primarily include coastal sage scrub, riparian scrub, nonnative grassland, ruderal vegetation, eucalyptus woodland, ornamental vegetation, and previously disturbed communities. Soil types mapped in the survey areas include the Hanford Association (0 to 5 percent slopes), Perkins-Rincon Association (0 to 15 percent slopes), and Altamont-Diablo Association (30 to 60 percent slopes, eroded).<sup>1</sup>

Prior to conducting focused surveys, LSA biologists reviewed appropriate literature to determine whether special-status plant species have been detected on or near the survey areas in the past. The

<sup>1</sup> LSA, 2007. Resource Management Plan, Puente Hills Landfill Native Habitat Preservation Authority. Appendix B: Soil Taxonomy and Analysis.



literature review included various documents prepared by LSA for the Habitat Authority (including previous survey documents), as well as the California Natural Diversity Database (CNDDDB), maintained by the California Department of Fish and Game (CDFG), and the California Native Plant Society's (CNPS) Online Inventory. The CNDDDB query included the *La Habra* and *Whittier, California* quadrangles, and the CNPS query included a 9-quad search of the surrounding quadrangles. LSA also reviewed the Resource Management Plan (RMP) prepared for the Habitat Authority in 2007, including the Sensitive Species Table in Appendix I, to further refine which special-status species might be present in the survey areas.

Based on the literature review, no special-status plants were found to have historic locations within the survey areas. However, nearby records for Plummer's mariposa lily (*Calochortus plummerae*) and Robinson's peppergrass (*Lepidium virginicum* var. *robinsonii*), both designated Special Plants by CDFG and on the CNPS List 1B, and Catalina mariposa lily (*Calochortus catalinae*), a CDFG Special Plant and CNPS List 4 species, suggest that there is high potential that these species could occur in the survey areas based on the presence of potentially suitable habitat. While the timing and methods of surveys focused on these three species, all vascular plant species observed in the survey areas were identified and recorded to document whether additional special-status species were present.

## METHODS

In 2008, the survey area included five project areas and totaled 113 acres. In 2009, the survey area included two large project areas, connected by a dirt road, which totaled 209 acres (Figure 2). LSA biologists surveyed all survey areas for each respective year on the following schedule:

LSA Biologists	Date	Time
Jim Harrison, Dan Rosie	April 10, 2008	7:00 a.m. to 4:30 p.m.
Dan Rosie, Jodi Ross	April 18, 2008	6:00 a.m. to 12:30 p.m.
Jim Harrison, Matthew Willis	June 4, 2008	6:30 a.m. to 3:30 p.m.
Jim Harrison, Dan Rosie	June 5, 2008	8:00 a.m. to 12:00 p.m.
Dan Rosie, Robert Steers	April 2, 2009	10:15 a.m. to 6:00 p.m.
Dan Rosie, Robert Steers	April 3, 2009	10:00 a.m. to 12:15 p.m.
Sarah Barrera, Robert Steers	June 5, 2009	11:00 a.m. to 3:30 p.m.
Sarah Barrera, Robert Steers	June 9, 2009	9:00 am to 12:00 p.m.

Botanical surveys were conducted in accordance with the current CNPS Botanical Survey Guidelines dated June 2, 2001. These surveys were floristic in nature, and every species noted in the field was identified to the extent necessary to determine whether it was a special-status plant species. In both years, the first survey was conducted in the middle of April to observe plants that emerge in early spring. The second botanical survey was performed during early June to observe plants that emerge during late spring.

The floristic surveys were conducted by walking transects throughout the survey areas. Transect widths varied from 10 to 100 feet and averaged approximately 50 feet, depending on visibility and habitat quality. Although the surveys were conducted during the expected flowering season in order to facilitate detection of these species, transects were walked slowly enough that they could have been

detected even in a preflowering or postflowering state. In addition, steep slopes inaccessible by foot were surveyed using binoculars from the most practical vantage points.

Precipitation during the study was 9.68 inches from September of 2007 to May 2008 (National Climate Data Center [NCDC] 2009) and 8.42 inches from September of 2008 to May 2009 in the nearby City of Whittier (NCDC 2009). Average precipitation for the City of Whittier is 14.05 inches from September through May, based on 59 years of data (Western Regional Climate Center 2009). Therefore precipitation was below average in both survey years. However, the majority of rainfall occurred between November and February in both years (data not shown), which led to widespread germination of native annual plants and bolting of perennial geophytes, like blue dicks (*Dichelostemma capitatum*) and blue-eyed grass (*Sisyrinchium bellum*). Furthermore, in 2009 the Habitat Authority ecologist confirmed that Plummer's mariposa lily was blooming at a nearby site during one of the survey visits.<sup>1</sup> Thus, it was concluded that these were adequate years and sampling dates to detect target special-status plant species.

A cumulative list of plant species identified during the April and June 2008 and 2009 surveys is found in attached Table A.

## RESULTS AND DISCUSSION

The target plant species were not detected in the survey areas during the April and June 2008 and 2009 focused surveys. A small stand (five individuals) of Southern California black walnut (*Juglans californica*) was found in the drainage that runs parallel to and east of Catalina Avenue, about 0.15 mile north of the entrance gate (see Figure 2). This small stand is a component of the riparian vegetation that occurs in the drainage. It does not constitute a California walnut woodland or walnut forest, which are vegetation types of interest to the CDFG. Southern California black walnut is on the CNPS 4.2 list. CNPS List 4 is only a "watch list," and species on this list do not appear on CNPS or CNDDB searches by USGS quads. This species has no State or federal status and is generally not afforded the same level of protection as species that are listed as threatened or endangered, but it is included in the Resource Management Plan for the preserve.

Historically, the survey areas have been heavily disturbed and much of the survey areas consist of nonnative vegetation. Patches of intact coastal sage scrub and other suitable habitat for special-status species do occur within the survey areas. However, the herbaceous component of these patches is dominated by nonnative species. Other areas in the survey areas that appeared relatively uninvaded did not contain special-status plants. The combination of historic disturbance and a high abundance of nonnative species likely preclude the existence for special-status species within the survey areas.

## CONCLUSIONS

Given the negative results of the focused surveys in 2008 and 2009, LSA concludes that Plummer's mariposa lily, Catalina mariposa lily, Robinson's peppergrass, or any other special-status plants do not occur within the 2008 or 2009 survey areas.

---

<sup>1</sup> Shannon Lucas, Habitat Authority Ecologist, pers. com., June 4, 2009.

If you have any questions or require additional information, please feel free to call Dan Rosie or me at (760) 931-5471 or e-mail one of us at Dan.Rosie@lsa-assoc.com or Robert.Steers@lsa-assoc.com, respectively.

Sincerely,

**LSA ASSOCIATES, INC.**

A handwritten signature in black ink that reads "Robert Steers". The signature is written in a cursive style with a large, looping initial "R".

Robert Steers, Ph.D.  
Biologist

Attachments: Figure 1: Project Location  
Figure 2: 2008 and 2009 Survey Areas and Results  
Table A: Plant Species Observed



## REFERENCES

- California Department of Fish and Game, Natural Heritage Division, Natural Diversity Database. 2008. RareFind Version 3.1.0. Records search executed April 8, 2008, covering the USGS 7.5-minute series topographic map, La Habra and Whittier, California quadrangles. Sacramento, California: The Resources Agency. Commercial version dated February 2, 2008.
- California Native Plant Society. 2001. Botanical Survey Guidelines. Revised June 2, 2001.
- . 2008. Inventory of Rare and Endangered Plants (online edition, v7-08b). California Native Plant Society. Sacramento, CA. Accessed on April 8, 2008. [<http://www.cnps.org/inventory>].
- Hickman, J.C. (ed.). 1996. The Jepson Manual: Higher Plants of California. University of California Press, Berkeley.
- Jepson Online Interchange for California Floristics. 2009. Accessed on June 23, 2009. [<http://ucjeps.berkeley.edu/interchange.html>]
- Ljubenkov, J.A.S., and T.S. Ross. 2002. An Annotated Checklist of the Vascular Plants of the Whittier Hills, Los Angeles County, California. *Crossosoma* 27(1).
- LSA Associates, Inc. 2006. Botanical Survey Report 2005. Prepared for the Puente Hills Landfill Native Habitat Preservation Authority. January 13 2006.
- LSA Associates, Inc. 2007. Resource Management Plan. Prepared for the Puente Hills Landfill Native Habitat Preservation Authority. July 2007.
- National Climatic Data Center. 2009. Annual Climatological Summary for Station: WHITTIER CITY YD FC106C, for 2007, 2008, and 2009. Website: <http://www.ncdc.noaa.gov/oa/climate/stationlocator.html> (Accessed on July 23, 2009).
- Western Regional Climate Center. 2009. Long Term Weather Summary for Station: WHITTIER CITY YD FC106C, Period of Record: 1/ 1/1949 to 12/31/2008. Website: <http://wrcc@dri.edu> (Accessed on July 23, 2009).









FIGURE 1

LSA


LEGEND

-  2009 Survey Area (approx. 209 acres)
-  2008 Survey Area (approx. 111 acres)



Puente Hills Landfill Native Habitat Preservation Authority  
 City of Whittier Oil Exploration  
 Project Location












LSA

FIGURE 2

LEGEND

- 
-  2009 Survey Areas (approx. 209 acres)
-  2008 Survey Areas (approx. 111 acres)
-  Trees mapped within Survey Area
-  Trees observed outside Survey Area (approx. 100)
-  Southern California Black Walnut (*Juglans californica*)



SOURCE: DigitalGlobe (April 2008)

Puente Hills Landfill Native Habitat Preservation Authority  
 City of Whittier Oil Exploration  
 2008 and 2009 Survey Areas and Results



## TABLE A

### VASCULAR PLANT SPECIES OBSERVED

The following vascular plant species were observed in the survey areas by various biologists during the course of on-site surveys in 2008 and 2009. Family- and species-level nomenclature follows the Jepson Manual. For species nomenclature only, current name changes on the Jepson Interchange are also used.

\*Introduced nonnative species

\*\*Special-status species

Scientific Name	Common Name
<b>ANGIOSPERMAE: DICOTYLEDONAE</b>	<b>DICOT FLOWERING PLANTS</b>
<b>Anacardiaceae</b>	<b>Sumac Family</b>
<i>Malosma laurina</i>	Laurel sumac
<i>Rhus integrifolia</i>	Lemonade berry
<i>Rhus ovata</i>	Sugar bush
* <i>Schinus molle</i>	Peruvian pepper tree
* <i>Schinus terebinthifolius</i>	Brazilian pepper tree
<i>Toxicodendron diversilobum</i>	Poison oak
<b>Apiaceae</b>	<b>Carrot Family</b>
* <i>Conium maculatum</i>	Poison hemlock
<i>Daucus pusillus</i>	American wild carrot
* <i>Foeniculum vulgare</i>	Sweet fennel
<b>Apocynaceae</b>	<b>Dogbane Family</b>
* <i>Vinca major</i>	Periwinkle
<b>Asclepiadaceae</b>	<b>Milweed Family</b>
<i>Asclepias californica</i>	California milkweed
<b>Asteraceae</b>	<b>Sunflower Family</b>
<i>Ambrosia acanthicarpa</i>	Annual sandbar
<i>Ambrosia psilostachya</i>	Western ragweed
<i>Artemisia californica</i>	California sagebrush
<i>Artemisia douglasiana</i>	California mugwort
<i>Baccharis emoryi</i>	Emory's baccharis
<i>Baccharis pilularis</i>	Coyote bush
<i>Baccharis salicifolia</i>	Mule fat
* <i>Carduus pycnocephalus</i>	Italian thistle
* <i>Centaurea melitensis</i>	Tocalote
* <i>Cirsium vulgare</i>	Bull thistle
<i>Corethrogyne filaginifolia</i>	California aster
<i>Deinandra fasciculata</i>	Fascicled tarweed
<i>Encelia californica</i>	California bush sunflower
<i>Gutierrezia californica</i>	California matchweed
* <i>Hedypnois cretica</i>	Crete hedypnois
<i>Helianthus annuus</i>	Annual sunflower
<i>Heterotheca grandiflora</i>	Telegraph weed

Scientific Name	Common Name
<i>Isocoma menziesii</i>	Coastal goldenbush
* <i>Lactuca serriola</i>	Prickly lettuce
<i>Corethrogyne filaginifolia</i>	Common sand aster
<i>Logfia filaginoides</i>	California filago
<i>Malacothrix saxatilis</i> var. <i>tenuifolia</i>	Cliff malacothrix
* <i>Picris echioides</i>	Bristly ox tongue
<i>Pseudognaphalium biolettii</i>	Two-color rabbit-tobacco
<i>Pseudognaphalium californicum</i>	California rabbit-tobacco
* <i>Pseudognaphalium luteoalbum</i>	Jersey cudweed
<i>Pseudognaphalium microcephalum</i>	San Diego rabbit-tobacco
* <i>Senecio vulgaris</i>	Common groundsel
* <i>Silybum marianum</i>	Milk thistle
* <i>Sonchus asper</i>	Common sow thistle
* <i>Sonchus oleraceus</i>	Common sow thistle
<i>Stephanomeria virgata</i>	Tall wreath plant
<i>Xanthium strumarium</i>	Cocklebur
<b>Brassicaceae</b>	<b>Mustard Family</b>
* <i>Brassica nigra</i>	Black mustard
* <i>Hirschfeldia incana</i>	Summer mustard
* <i>Raphanus sativus</i>	Wild radish
* <i>Sisymbrium</i> sp.	Mustard
* <i>Sisymbrium irio</i>	London rocket
<b>Cactaceae</b>	<b>Cactus Family</b>
<i>Opuntia littoralis</i>	Coastal prickly pear
<b>Caprifoliaceae</b>	<b>Honeysuckle Family</b>
<i>Sambucus mexicana</i>	Mexican elderberry
<b>Caryophyllaceae</b>	<b>Pink Family</b>
<i>Stellaria media</i>	Chickweed
<b>Chenopodiaceae</b>	<b>Goosefoot Family</b>
* <i>Atriplex semibaccata</i>	Australian saltbush
* <i>Chenopodium album</i>	Lamb's quarters
<i>Chenopodium berlandieri</i>	Pitseed goosefoot
* <i>Chenopodium murale</i>	Nettle leaf goosefoot
* <i>Salsola tragus</i>	Russian thistle
<b>Convolvulaceae</b>	<b>Morning-Glory Family</b>
<i>Calystegia macrostegia</i>	Morning-glory
<b>Crassulaceae</b>	<b>Stonecrop Family</b>
<i>Crassula connata</i>	Sand pygmy-stonecrop
<b>Cucurbitaceae</b>	<b>Gourd Family</b>
<i>Cucurbita foetidissima</i>	Coyote melon
<i>Marah macrocarpus</i>	Man root
<b>Euphorbiaceae</b>	<b>Spurge Family</b>
<i>Chamaesyce albomarginata</i>	Rattlesnake weed
<i>Chamaesyce</i> sp.	Spurge
* <i>Ricinis communis</i>	Castor bean
<b>Fabaceae</b>	<b>Legume Family</b>
* <i>Acacia cyclops</i>	Coastal wattle
* <i>Acacia longifolia</i>	Golden wattle
<i>Lotus salsuginosus</i> var. <i>salsuginosus</i>	Coastal lotus

Scientific Name	Common Name
<i>Lotus scoparius</i>	Deer weed
<i>Lupinus microcarpus</i> var. <i>densiflorus</i>	Dense-flowered chick lupine
<i>Lupinus succulentus</i>	Arroyo lupine
* <i>Medicago polymorpha</i>	Bur clover
* <i>Melilotus indica</i>	Yellow sweet clover
<b>Fagaceae</b>	<b>Oak Family</b>
<i>Quercus agrifolia</i>	Coast live oak
<b>Geraniaceae</b>	<b>Geranium Family</b>
* <i>Erodium cicutarium</i>	Red-stemmed filaree
* <i>Erodium moschatum</i>	White-stemmed filaree
<b>Grossulariaceae</b>	<b>Gooseberry Family</b>
<i>Ribes speciosum</i>	Fuchsia flowered gooseberry
<b>Hydrophyllaceae</b>	<b>Waterleaf Family</b>
<i>Emmenanthe penduliflora</i>	Whispering bells
<i>Eucrypta chrysanthemifolia</i>	Common eucrypta
<i>Phacelia cicutaria</i> var. <i>hispida</i>	Caterpillar phacelia
<i>Phacelia minor</i>	California bluebell
<i>Phacelia parryi</i>	Parry's phacelia
<i>Phacelia ramosissima</i>	Branching phacelia
<i>Phacelia tanacetifolia</i>	Tansy leafed phacelia
<i>Pholistoma auritum</i> var. <i>auritum</i>	Blue fiesta flower
<b>Juglandaceae</b>	<b>Walnut Family</b>
** <i>Juglans californica</i>	Southern California walnut
<b>Lamiaceae</b>	<b>Mint Family</b>
* <i>Marrubium vulgare</i>	Horehound
<i>Salvia apiana</i>	White sage
<i>Salvia leucophylla</i> X <i>apiana</i>	White/purple sage hybrid
<i>Salvia leucophylla</i>	Purple sage
<i>Salvia mellifera</i>	Black sage
<b>Malvaceae</b>	<b>Mallow Family</b>
<i>Malacothamnus fasciculatus</i>	Lax-flowered mallow
* <i>Malva parviflora</i>	Cheeseweed
* <i>Malva sylvestris</i>	High mallow
<b>Myrtaceae</b>	<b>Myrtle Family</b>
* <i>Eucalyptus</i> sp.	Gum
<b>Nyctaginaceae</b>	<b>Four-O'Clock Family</b>
<i>Mirabilis laevis</i>	Wishbone bush
<b>Onagraceae</b>	<b>Evening Primrose Family</b>
<i>Camissonia californica</i>	California suncup
<i>Clarkia bota</i>	Botta's clarkia
<b>Oxalidaceae</b>	<b>Oxalis Family</b>
* <i>Oxalis pes-caprae</i>	Bermuda buttercup
<b>Platanaceae</b>	<b>Sycamore Family</b>
<i>Platanus racemosa</i>	California sycamore
<b>Polygonaceae</b>	<b>Buckwheat Family</b>
<i>Eriogonum fasciculatum</i>	California buckwheat
* <i>Rumex crispus</i>	Curly dock
<b>Portulacaceae</b>	<b>Purselane Family</b>
<i>Claytonia perfoliata</i>	Miner's lettuce



Scientific Name	Common Name
<b>Primulaceae</b>	<b>Primrose Family</b>
* <i>Anagallis arvensis</i>	Scarlet pimpernel
<b>Rosaceae</b>	<b>Rose Family</b>
<i>Heteromeles arbutifolia</i>	Toyon
<i>Prunus ilicifolia</i>	Hollyleaf cherry
<b>Rubiaceae</b>	<b>Madder Family</b>
<i>Galium angustifolium</i>	Chaparral bedstraw
<i>Galium aparine</i>	Common bedstraw
<b>Salicaceae</b>	<b>Willow Family</b>
<i>Salix exigua</i>	Narrowleaf willow
<i>Salix gooddingii</i>	Black willow
<i>Salix laevigata</i>	Red willow
<i>Salix lasiolepis</i>	Arroyo willow
<b>Scrophulariaceae</b>	<b>Figwort Family</b>
<i>Keckiella cordifolia</i>	Heart-leaved penstemon
<i>Mimulus aurantiacus</i>	Orange bush monkey flower
<i>Scrophularia californica</i>	California figwort
* <i>Verbascum virgatum</i>	Wandmullein
<b>Solanaceae</b>	<b>Nightshade Family</b>
* <i>Nicotiana glauca</i>	Tree tobacco
* <i>Solanum americanum</i>	White nightshade
<i>Solanum douglasii</i>	Douglas' nightshade
<b>Tamaricaceae</b>	<b>Tamarisk Family</b>
* <i>Tamarix ramosissima</i>	Mediterranean tamarisk
<b>Tropaeolaceae</b>	<b>Nightshade Family</b>
* <i>Tropaeolum majus</i>	Garden nasturtium
<b>Ulmaceae</b>	<b>Elm Family</b>
* <i>Ulmus parvifolia</i>	Chinese elm
<b>Urticaceae</b>	<b>Nettle Family</b>
<i>Urtica dioica</i> ssp. <i>holosericea</i>	Hoary nettle
* <i>Urtica urens</i>	Dwarf nettle
<i>Parietaria hespera</i>	Western pellitory
<b>Verbenaceae</b>	<b>Vervain Family</b>
<i>Verbena lasiostachys</i>	Western verbena
<b>ANGIOSPERMAE: MONOCOTYLEDONAE</b>	<b>MONOCOT FLOWERING PLANTS</b>
<b>Arecaceae</b>	<b>Palm Family</b>
* <i>Washingtonia robusta</i>	Mexican fan palm
<b>Iridaceae</b>	<b>Iris Family</b>
<i>Sisyrinchium bellum</i>	Blue eyed grass
<b>Liliaceae</b>	<b>Lily Family</b>
<i>Dichelostemma capitatum</i>	Blue dicks
<b>Poaceae</b>	<b>Grass Family</b>
* <i>Avena barbata</i>	Slender wild oat
* <i>Avena fatua</i>	Wild oat
* <i>Bromus diandrus</i>	Ripgut brome
* <i>Bromus hordeaceus</i>	Soft chess
* <i>Bromus madritensis</i> ssp. <i>rubens</i>	Red brome
* <i>Cortaderia jubata</i>	Pampas grass
* <i>Cynodon dactylon</i>	Bermuda grass

<b>Scientific Name</b>	<b>Common Name</b>
* <i>Hordeum murinum</i> ssp. <i>leporinum</i>	Foxtail barley
<i>Leymus condensatus</i>	Giant wildrye
<i>Leymus triticoides</i>	Creeping wild rye
* <i>Lolium multiflorum</i>	Italian ryegrass
<i>Nassella pulchra</i>	Purple needlegrass
<i>Nassella lepida</i>	Foothill needlegrass
* <i>Pennisetum setaceum</i>	African fountain grass
* <i>Piptatherum miliaceum</i>	Smilo grass
* <i>Schismus barbatus</i>	Schismus
* <i>Vulpia myuros</i>	Foxtail fescue



LSA ASSOCIATES, INC.  
PACIFIC CENTER  
703 PALOMAR AIRPORT RD., SUITE 260  
CARLSBAD, CALIFORNIA 92011

760.931.5471 TEL  
760.918.2458  
FAX

BERKELEY  
COLMA  
FORT COLLINS

IRVINE  
PALM SPRINGS  
POINT RICHMOND

RIVERSIDE  
ROCKLIN  
SAN LUIS OBISPO

August 27, 2008

Ms. Sandra Marquez  
United States Fish and Wildlife Service  
Carlsbad Field Office  
6010 Hidden Valley Road  
Carlsbad, CA 92011

Permit Biologist  
Nongame Unit, Wildlife Branch  
California Department of Fish and Game  
1416 Ninth Street, 12th Floor  
Sacramento, CA 95814

Subject: Least Bell's Vireo and Coastal California Gnatcatcher Survey Results, Puente Hills  
Landfill Native Habitat Preservation Authority Lands, City of Whittier, Los Angeles  
County, California (LSA Project No. PUE0801)

Dear Ms. Marquez and Permit Biologist:

This letter report documents the results of protocol surveys for the least Bell's vireo (*Vireo bellii pusillus*) and coastal California gnatcatcher (*Polioptila californica californica*) conducted by LSA Associates, Inc. (LSA). Surveys were conducted at five potential oil extraction sites within Puente Hills Landfill Native Habitat Preservation Authority (Habitat Authority) Lands in the City of Whittier, Los Angeles County, California (Figure 1, attached). The surveys were requested by United States Fish and Wildlife Service (USFWS) biologist Ken Corey during an on-site visit March 6, 2008. The least Bell's vireo is a USFWS federally endangered species and a California Department of Fish and Game (CDFG) endangered species. The coastal California gnatcatcher is a USFWS federally threatened species.

No least Bell's vireos or coastal California gnatcatchers were detected during protocol surveys, but one coastal California gnatcatcher was observed in the study area by a Habitat Authority employee.

## STUDY AREA

The project area includes five sites under investigation for potential oil exploration activities within lands managed by the Habitat Authority. Combined, the sites total approximately 113 acres and are located within Sections 22, 23, and 26 of Township 2 South, Range 11 West, as shown on the United States Geological Survey (USGS) 7.5-minute *Whittier* and *La Habra, California* quadrangles (Figure 1).

The five project sites are characterized by portions of the Arroyo Pescadero and La Canada Verde drainages, adjacent hillsides, and access roads. Elevation at the site ranges from approximately 300 to 1,000 feet above mean sea level. Vegetation communities within the project sites primarily include coastal sage scrub, riparian scrub, nonnative grassland, ruderal vegetation, eucalyptus woodland, ornamental vegetation, and other previously disturbed areas.



## METHODS

Richard Erickson conducted eight protocol least Bell’s vireo and coastal California gnatcatcher surveys from April 21 to July 9, 2008. During each of the surveys, the biologist walked slowly along the edge of and, when appropriate, through riparian and coastal sage scrub habitat, listening for least Bell’s vireos and coastal California gnatcatchers. A taped recording of a coastal California gnatcatcher was played periodically along the survey route during all of the surveys. The surveying biologist, with the aid of binoculars for viewing wildlife species, waited for several minutes after each playing to look and listen for both least Bell’s vireos and coastal California gnatcatchers.

**Table A: Survey Schedule and Conditions**

Date 2008	Time	Weather	Surveyor
April 21	0640–1040	Clear, cool, light wind	RE
May 1	0600–0940	Nearly complete marine layer, cool, calm	RE
May 12	0555–0920	Complete marine layer with some heavy mist, mild, calm	RE
May 22	0540–0935	Edge of complete marine layer, mild, light wind	RE
June 3	0515–0915	Complete marine layer, cool, calm	RE
June 13	0535–0930	Edge of complete marine layer, mild, light wind	RE
June 27	0610–0955	Complete marine layer, mild, calm	RE
July 9	0540–0945	Complete heavy marine layer, mild, calm	RE

Surveyor: Richard Erickson.

Surveys were conducted pursuant to Federal Fish and Wildlife Permit TE777965-8 (April 8, 2008–April 7, 2012) and a temporary authorization from the CDFG (May 12, 2003–March 31, 2007; renewal request submitted March 26, 2007, extending coverage indefinitely) in lieu of a Memorandum of Understanding between LSA and CDFG.

## RESULTS

No least Bell’s vireos or coastal California gnatcatchers were found during the surveys. However, a single coastal California gnatcatcher was seen in the study area (Figure 2) by Habitat Authority employee David Moskovitz on June 9 and 10. A search of the same area by Mr. Moskovitz and Mr. Erickson on June 13 was unsuccessful. The bird was most likely a recently fledged juvenile dispersing from its natal territory.


The brown-headed cowbird (*Molothrus ater*) – a brood parasite of least Bell’s vireos, coastal California gnatcatchers, and other passerines – was detected on two of the first three surveys, with a maximum of three birds noted on May 1.

A complete list of the animals detected on these surveys is included in Appendix A (attached).

If you have any questions, please contact me at (949) 553-0666.

Sincerely,


**LSA ASSOCIATES, INC.**



Richard Erickson  
Associate/Biologist

Attachments: Figure 1: Project Location  
Figure 2: Survey Areas  
Appendix A: Animal Species Observed  
Appendix B: California Native Species Field Survey Summary Form

**I CERTIFY THAT THE INFORMATION IN THIS SURVEY REPORT AND ATTACHED EXHIBITS FULLY AND ACCURATELY REPRESENTS MY WORK:**

<b>SURVEYOR:</b>	<b>PERMIT NUMBER</b>	<b>DATE:</b>
 Richard Erickson	TE-777965-8	August 7, 2008







FIGURE 1

LSA

LEGEND

Project Sites (Survey Area)



Puente Hills Landfill Native Habitat Preservation Authority  
 Focused Surveys for City of Whittier Oil Exploration  
 Project Location

SOURCE: USGS 7.5' QUAD - WHITTIER (81) AND LA HABRA (81); CALIF.




E:\PUE0801\GIS\Fig1\_location.mxd (08/05/08)






LSA

LEGEND

-  Coastal California gnatcatcher observed by Habitat Authority employee D. Moskovitz June 9 & 10, 2008
-  Survey Areas (Acres) - all areas are potentially suitable for dispersing juvenile coastal California gnatcatcher
-  Potential least Bell's vireo nesting habitat near survey areas



SOURCE: Airphoto USA (June 2007); Matrix Oil Corp (June 2008)

I:\PUE0801\GIS\Fig2\_Project\_Sites\_w\_CAGN.mxd (08/25/08)

FIGURE 2

Puente Hills Landfill Native Habitat Preservation Authority  
 Focused Surveys for City of Whittier Oil Exploration  
 Survey Areas



## **APPENDIX A**

### **ANIMAL SPECIES OBSERVED**

## APPENDIX A

### ANIMAL SPECIES OBSERVED

This is a list of the conspicuous aerial insects, amphibians, reptiles, birds, and mammals noted in the study area by LSA biologists. Presence may be noted if a species is seen or heard, or identified by the presence of tracks, scat, or other signs.

\* Species not native to the study area

#### ANISOPTERA

##### Libellulidae

*Pantala hymenaea*

#### LEPIDOPTERA

##### Pieridae

*Pontia protodice*

\* *Pieris rapae*

*Anthocharis sara*

*Colias eurytheme*

##### Lycaenidae

*Leptotes marina*

##### Nymphalidae

*Vanessa cardui*

*Vanessa anabella*

*Vanessa atalanta*

#### AMPHIBIA

##### Hylidae

*Pseudacris hypochondriaca*

#### REPTILIA

##### Phrynosomatidae

*Sceloporus occidentalis*

#### TYPICAL DRAGONFLIES

##### Cruisers, Emeralds, Baskettails, and Skimmers

Spot-winged glider

#### BUTTERFLIES

##### Whites and Sulphurs

Checkered white

Cabbage white

Sara orangetip

Orange sulphur

##### Gossamer-Wing Butterflies

Marine blue

##### Brush-Footed Butterflies

Painted lady

West coast lady

Red admiral

#### AMPHIBIANS

##### Treefrogs and Relatives

Baja California treefrog

#### REPTILES

##### Phrynosomatid Lizards

Western fence lizard



**Teiidae**

*Aspidoscelis tigris*

**AVES**

**Odontophoridae**

*Callipepla californica*

**Phalacrocoracidae**

*Phalacrocorax auritus*

**Ardeidae**

*Ardea herodias*

**Cathartidae**

*Cathartes aura*

**Accipitridae**

*Accipiter cooperii*

*Buteo lineatus*

*Buteo jamaicensis*

**Falconidae**

*Falco sparverius*

**Laridae**

*Larus californicus*

*Larus occidentalis*

**Columbidae**

\* *Columba livia*

*Patagioenas fasciata*

*Zenaida macroura*

**Psittacidae**

\* *Brotogeris chiriri*

**Strigidae**

*Bubo virginianus*

**Apodidae**

*Aeronautes saxatilis*

**Trochilidae**

*Archilochus alexandri*

*Calypte anna*

**Whiptails and Relatives**

Western whiptail

**BIRDS**

**New World Quail**

California quail

**Cormorants**

Double-crested cormorant

**Herons, Bitterns, and Allies**

Great blue heron

**New World Vultures**

Turkey vulture

**Hawks, Kites, Eagles, and Allies**

Cooper's hawk

Red-shouldered hawk

Red-tailed hawk

**Caracaras and Falcons**

American kestrel

**Gulls, Terns, and Skimmers**

California gull

Western gull

**Pigeons and Doves**

Rock (Feral) pigeon

Band-tailed pigeon

Mourning dove

**Lories, Parakeets, Macaws, and Parrots**

Yellow-chevroned parakeet

**Typical Owls**

Great horned owl

**Swifts**

White-throated swift

**Hummingbirds**

Black-chinned hummingbird

Anna's hummingbird

*Calypte costae*  
*Selasphorus rufus/sasin*

Costa's hummingbird  
Rufous/Allen's hummingbird

**Picidae**

*Melanerpes formicivorus*  
*Sphyrapicus* sp.  
*Picoides nuttallii*  
*Picoides pubescens*  
*Colaptes auratus*

**Woodpeckers and Allies**

Acorn woodpecker  
Sapsucker sp.  
Nuttall's woodpecker  
Downy woodpecker  
Northern flicker

**Tyrannidae**

*Empidonax difficilis*  
*Sayornis nigricans*  
*Myiarchus cinerascens*  
*Tyrannus vociferans*  
*Tyrannus verticalis*

**Tyrant Flycatchers**

Pacific-slope flycatcher  
Black phoebe  
Ash-throated flycatcher  
Cassin's kingbird  
Western kingbird

**Vireonidae**

*Vireo huttoni*  
*Vireo gilvus*

**Vireos**

Hutton's vireo  
Warbling vireo

**Corvidae**

*Aphelocoma californica*  
*Corvus brachyrhynchos*  
*Corvus corax*

**Crows and Jays**

Western scrub-jay  
American crow  
Common raven

**Hirundinidae**

*Stelgidopteryx serripennis*  
*Petrochelidon pyrrhonota*  
*Hirundo rustica*

**Swallows**

Northern rough-winged swallow  
Cliff swallow  
Barn swallow

**Aegithalidae**

*Psaltriparus minimus*

**Long-Tailed Tits and Bushtits**

Bushtit

**Troglodytidae**

*Thryomanes bewickii*  
*Troglodytes aedon*

**Wrens**

Bewick's wren  
House wren

**Turdidae**

*Sialia mexicana*  
*Catharus ustulatus*  
*Turdus migratorius*

**Thrushes**

Western bluebird  
Swainson's thrush  
American robin

**Timaliidae**

*Chamaea fasciata*

**Babblers**

Wrentit

**Mimidae**

*Mimus polyglottos*  
*Toxostoma redivivum*

**Bombycillidae**

*Bombycilla cedrorum*

**Ptilonotidae**

*Phainopepla nitens*

**Parulidae**

*Vermivora celata*  
*Vermivora ruficapilla*  
*Dendroica petechia*  
*Dendroica coronata*  
*Dendroica nigrescens*  
*Dendroica townsendi*  
*Dendroica occidentalis*  
*Geothlypis trichas*  
*Wilsonia pusilla*  
*Icteria virens*

**Thraupidae**

*Piranga ludoviciana*

**Emberizidae**

*Pipilo maculatus*  
*Pipilo crissalis*  
*Aimophila ruficeps*  
*Spizella passerina*  
*Melospiza melodia*  
*Zonotrichia atricapilla*

**Cardinalidae**

*Pheucticus melanocephalus*  
*Passerina caerulea*  
*Passerina amoena*

**Icteridae**

*Agelaius phoeniceus*  
*Molothrus ater*  
*Icterus cucullatus*  
*Icterus bullockii*

**Fringillidae**

*Carpodacus mexicanus*

**Mockingbirds and Thrashers**

Northern mockingbird  
California thrasher

**Waxwings**

Cedar waxwing

**Silky-flycatchers**

Phainopepla

**Wood Warblers**

Orange-crowned warbler  
Nashville warbler  
Yellow warbler  
Yellow-rumped warbler  
Black-throated gray warbler  
Townsend's warbler  
Hermit warbler  
Common yellowthroat  
Wilson's warbler  
Yellow-breasted chat

**Tanagers**

Western tanager

**Emberizids**

Spotted towhee  
California towhee  
Rufous-crowned sparrow  
Chipping sparrow  
Song sparrow  
Golden-crowned sparrow

**Cardinals, Saltators, and Allies**

Black-headed grosbeak  
Blue grosbeak  
Lazuli bunting

**Blackbirds**

Red-winged blackbird  
Brown-headed cowbird  
Hooded oriole  
Bullock's oriole

**Fringilline and Cardueline Finches and Allies**

House finch



*Carduelis psaltria*  
*Carduelis tristis*

Lesser goldfinch  
American goldfinch

**Passeridae**

\* *Passer domesticus*

**Old World Sparrows**

House sparrow

**Estrildidae**

\* *Lonchura punctulata*

**Estrildid Finches**

Nutmeg mannikin

**MAMMALIA**

**MAMMALS**

**Sciuridae**

\* *Sciurus niger*  
*Spermophilus beecheyi*

**Squirrels, Chipmunks, and Marmots**

Eastern fox squirrel  
California ground squirrel

**Geomyidae**

*Thomomys bottae*

**Pocket Gophers**

Botta's pocket gopher

**Cricetidae**

*Neotoma lepida*  
*Neotoma macrotis*

**Hamsters, Voles, Lemmings, and New  
World Rats and Mice**

Desert woodrat  
Big-eared woodrat

**Leporidae**

*Sylvilagus audubonii*

**Rabbits and Hares**

Audubon's cottontail

**Canidae**

*Canis latrans*

**Foxes, Wolves, and Allies**

Coyote

**Cervidae**

*Odocoileus hemionus*

**Deer, Elk, and Allies**

Mule deer

**Taxonomy and nomenclature are based on the following.**

Damselflies and dragonflies: Manolis, T. (2003, Dragonflies and Damselflies of California, University of California Press, Berkeley).

Butterflies: North American Butterfly Association (2001, NABA checklist and English Names of North American Butterflies, Second Edition, North American Butterfly Association, Morristown, New Jersey).

Amphibians and reptiles: Crother, B.I. ed. (2008. Scientific and Standard English Names of Amphibians and Reptiles of North America North of Mexico. *Herpetological Circular* 37) for species taxonomy and nomenclature; Stebbins, R.C. (2003, A Field Guide to Western Reptiles and Amphibians, third edition, Houghton Mifflin, Boston) for sequence and higher order taxonomy.

Birds: American Ornithologists' Union (1998, The A.O.U. Checklist of North American Birds, Seventh Edition, American Ornithologists' Union, Washington D.C.; and 2000, 2002, 2003, 2004, 2005, 2006, and 2007 supplements; see <http://aou.org.whsites.net/checklist/index.php3>).

Mammals: Wilson, D.E., and D.M. Reeder, eds. (2005. Mammal Species of the World, 3rd ed. Johns Hopkins University Press, Baltimore, Maryland; see <http://nmmhgoph.si.edu/msw/>).

**APPENDIX B**

**CALIFORNIA NATIVE SPECIES FIELD SURVEY SUMMARY FORM**



Mail to:  
 California Natural Diversity Database  
 Department of Fish and Game  
 1807 13<sup>th</sup> Street, Suite 202  
 Sacramento, CA 95811  
 Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov

For Office Use Only

Source Code \_\_\_\_\_ Quad Code \_\_\_\_\_  
 Elm Code \_\_\_\_\_ Occ. No. \_\_\_\_\_  
 EO Index No. \_\_\_\_\_ Map Index No. \_\_\_\_\_

Date of Field Work (mm/dd/yyyy): 4/21-7/9/2008

Reset

California Native Species Field Survey Form

Send Form

**Scientific Name:** Vireo bellii pusillus

**Common Name:** Least Bell's vireo

**Species Found?**  Yes  No not present?  
If not, why?

**Reporter:** Richard A. Erickson

**Address:** LSA Associates 20 Executive Park, Suite 200, Irvine, CA 92614

**E-mail Address:** richard.erickson@LSA-assoc.com

**Phone:** 949 553-0666

Total No. Individuals \_\_\_\_\_ Subsequent Visit?  yes  no

Is this an existing NDDDB occurrence? \_\_\_\_\_  no  unk.  
Yes, Occ. #

Collection? If yes: \_\_\_\_\_  
Number Museum / Herbarium

<p><b>Plant Information</b></p> <p>Phenology: _____% vegetative _____% flowering _____% fruiting</p>	<p><b>Animal Information</b></p> <p># adults <input type="checkbox"/> breeding                  # juveniles <input type="checkbox"/> wintering                  # larvae <input type="checkbox"/> burrow site                  # egg masses <input type="checkbox"/> rookery                  # unknown <input type="checkbox"/> nesting other</p>
--	--

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

Puente Hills, Whittier Puente Hills Landfill Native

County: Los Angeles Landowner / Mgr.: Habitat Preservation Authority

Quad Name: Whittier and La Habra Elevation: 300-1000 ft

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  W  
 Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  W  
 GPS Make & Model \_\_\_\_\_

**DATUM:** NAD27  NAD83  WGS84

Horizontal Accuracy \_\_\_\_\_ meters/feet

**Coordinate System:** UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)

Coordinates: \_\_\_\_\_

**Habitat Description (plant communities, dominants, associates, substrates/soils, aspects/slope):**

coastal sage scrub, ruderal grassland, riparian scrub & woodland, eucalyptus woodland

Other rare taxa seen at THIS site on THIS date: Dendroica petechia, Icteria virens  
Neotoma lepida intermedia

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor

Immediate AND surrounding land use: native habitat preserve

Visible disturbances: invasive exotic vegetation, extensive road system

Threats: oil extraction?

Comments: \_\_\_\_\_

<p><b>Determination:</b> (check one or more, and fill in blanks)</p> <p><input type="checkbox"/> Keyed (cite reference): _____</p> <p><input type="checkbox"/> Compared with specimen housed at: _____</p> <p><input type="checkbox"/> Compared with photo / drawing in: _____</p> <p><input type="checkbox"/> By another person (name): _____</p> <p><input checked="" type="checkbox"/> Other: _____</p>	<p><b>Photographs:</b> (check one or more)</p> <table border="1"> <tr> <td>Slide</td> <td>Print</td> <td>Digital</td> </tr> <tr> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> <tr> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> <tr> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> </table> <p>Plant / animal                  Habitat                  Diagnostic feature</p> <p>May we obtain duplicates at our expense? yes <input type="checkbox"/> no <input type="checkbox"/></p>	Slide	Print	Digital	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Slide	Print	Digital											
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>											
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>											
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>											

Mail to:  
 California Natural Diversity Database  
 Department of Fish and Game  
 1807 13<sup>th</sup> Street, Suite 202  
 Sacramento, CA 95811

Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov

**For Office Use Only**

Source Code \_\_\_\_\_ Quad Code \_\_\_\_\_  
 Elm Code \_\_\_\_\_ Occ. No. \_\_\_\_\_  
 EO Index No. \_\_\_\_\_ Map Index No. \_\_\_\_\_

**Date of Field Work (mm/dd/yyyy):** 4/21-7/9/2008

**Reset**

**California Native Species Field Survey Form**

**Send Form**

**Scientific Name:** Poliophtila californica californica

**Common Name:** coastal California gnatcatcher

**Species Found?**  Yes  No \*see under Site Information below  
If not, why?

Total No. Individuals \_\_\_\_\_ Subsequent Visit?  yes  no

**Is this an existing NDDDB occurrence?**  no  unk.  
Yes, Occ. #

Collection? If yes: \_\_\_\_\_  
Number Museum / Herbarium

**Reporter:** Richard A. Erickson

**Address:** LSA Associates 20 Executive Park, Suite 200, Irvine, CA 92614

**E-mail Address:** richard.erickson@LSA-assoc.com

**Phone:** 949 553-0666

Plant Information	Animal Information
Phenology: _____% vegetative _____% flowering _____% fruiting	# adults _____ # juveniles _____ # larvae _____ # egg masses _____ # unknown _____
	<input type="checkbox"/> breeding <input type="checkbox"/> wintering <input type="checkbox"/> burrow site <input type="checkbox"/> rookery <input type="checkbox"/> nesting <input type="checkbox"/> other

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

Puente Hills, Whittier Puente Hills Landfill Native

County: Los Angeles Landowner / Mgr.: Habitat Preservation Authority

Quad Name: Whittier and La Habra Elevation: 300-1000 ft

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  U  
 Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  U  
 GPS Make & Model \_\_\_\_\_

**DATUM:** NAD27  NAD83  WGS84

Horizontal Accuracy \_\_\_\_\_ meters/feet

**Coordinate System:** UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)

**Coordinates:** \_\_\_\_\_

**Habitat Description (plant communities, dominants, associates, substrates/soils, aspects/slope):**

coastal sage scrub, ruderal grassland, riparian scrub & woodland, eucalyptus woodland

Other rare taxa seen at THIS site on THIS date: Dendroica petechia, Icteria virens  
 (separate form preferred) Neotoma lepida intermedia

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor

Immediate AND surrounding land use: native habitat preserve

Visible disturbances: invasive exotic vegetation, extensive road system

Threats: oil extraction?

Comments: one coastal California gnatcatcher was observed here by PHLNHPA employee David Moskowitz June 9-10, 2008; it was presumably a dispersing juvenile.

Determination: (check one or more, and fill in blanks)	Photographs: (check one or more)
<input type="checkbox"/> Keyed (cite reference): _____	Plant / animal <input type="checkbox"/> Slide <input type="checkbox"/> Print <input type="checkbox"/> Digital <input type="checkbox"/>
<input type="checkbox"/> Compared with specimen housed at: _____	Habitat <input type="checkbox"/> Slide <input type="checkbox"/> Print <input type="checkbox"/> Digital <input type="checkbox"/>
<input type="checkbox"/> Compared with photo / drawing in: _____	Diagnostic feature <input type="checkbox"/> Slide <input type="checkbox"/> Print <input type="checkbox"/> Digital <input type="checkbox"/>
<input type="checkbox"/> By another person (name): _____	
<input checked="" type="checkbox"/> Other: _____	May we obtain duplicates at our expense? yes <input type="checkbox"/> no <input type="checkbox"/>

*For Office Use Only*

Source Code \_\_\_\_\_ Quad Code \_\_\_\_\_  
 Elm Code \_\_\_\_\_ Occ. No. \_\_\_\_\_  
 EO Index No. \_\_\_\_\_ Map Index No. \_\_\_\_\_

Date of Field Work (mm/dd/yyyy): 5/1-12/2008

Reset

## California Native Species Field Survey Form

Send Form

**Scientific Name:** Dendroica petechia

**Common Name:** yellow warbler

**Species Found?**  Yes  No If not, why?

Total No. Individuals 2 or 3 Subsequent Visit?  yes  no

Is this an existing NDDDB occurrence?  no  unk. Yes, Occ. #

Collection? If yes: \_\_\_\_\_  
Number Museum / Herbarium

**Reporter:** Richard A. Erickson

**Address:** LSA Associates 20 Executive Park, Suite 200, Irvine, CA 92614

**E-mail Address:** richard.erickson@LSA-assoc.com

**Phone:** 949 553-0666

Plant Information	Animal Information
Phenology: _____% vegetative _____% flowering _____% fruiting	<u>2 or 3</u> # adults _____ # juveniles _____ # larvae _____ # egg masses _____ # unknown _____ <input type="checkbox"/> breeding <input type="checkbox"/> wintering <input type="checkbox"/> burrow site <input type="checkbox"/> rookery <input type="checkbox"/> nesting <input checked="" type="checkbox"/> other <u>migrating through</u>

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

Puente Hills, Whittier Puente Hills Landfill Native

County: Los Angeles Landowner / Mgr.: Habitat Preservation Authority

Quad Name: Whittier and La Habra Elevation: 300-1000 ft

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  GPS Make & Model \_\_\_\_\_

**DATUM:** NAD27  NAD83  WGS84  Horizontal Accuracy \_\_\_\_\_ meters/feet

Coordinate System: UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)

Coordinates: \_\_\_\_\_

**Habitat Description (plant communities, dominants, associates, substrates/soils, aspects/slope):**

coastal slope scrub, ruderal grassland, riparian scrub & woodland, eucalyptus woodland

Other rare taxa seen at THIS site on THIS date:  
 (separate form preferred)

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor

Immediate AND surrounding land use: native habitat preserve

Visible disturbances: invasive exotic vegetation, extensive road system

Threats: oil extraction?

Comments:

<p><b>Determination:</b> (check one or more, and fill in blanks)</p> <p><input type="checkbox"/> Keyed (cite reference): _____</p> <p><input type="checkbox"/> Compared with specimen housed at: _____</p> <p><input type="checkbox"/> Compared with photo / drawing in: _____</p> <p><input type="checkbox"/> By another person (name): _____</p> <p><input checked="" type="checkbox"/> Other: _____</p>	<p><b>Photographs:</b> (check one or more)</p> <table style="width: 100%;"> <tr> <td>Slide</td> <td>Print</td> <td>Digital</td> </tr> <tr> <td>Plant / animal</td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> <tr> <td>Habitat</td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> <tr> <td>Diagnostic feature</td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> </table> <p>May we obtain duplicates at our expense? yes <input type="checkbox"/> no <input type="checkbox"/></p>	Slide	Print	Digital	Plant / animal	<input type="checkbox"/>	<input type="checkbox"/>	Habitat	<input type="checkbox"/>	<input type="checkbox"/>	Diagnostic feature	<input type="checkbox"/>	<input type="checkbox"/>
Slide	Print	Digital											
Plant / animal	<input type="checkbox"/>	<input type="checkbox"/>											
Habitat	<input type="checkbox"/>	<input type="checkbox"/>											
Diagnostic feature	<input type="checkbox"/>	<input type="checkbox"/>											



California Natural Diversity Database  
Department of Fish and Game  
1807 13<sup>th</sup> Street, Suite 202  
Sacramento, CA 95811

Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov

## For Office Use Only

Source Code \_\_\_\_\_ Quad Code \_\_\_\_\_  
Elm Code \_\_\_\_\_ Occ. No. \_\_\_\_\_  
EO Index No. \_\_\_\_\_ Map Index No. \_\_\_\_\_

Date of Field Work (mm/dd/yyyy): 6/13-7/9/2008

Reset

## California Native Species Field Survey Form

Send Form

Scientific Name: <u>Icteria virens</u>	
Common Name: <u>yellow-breasted chat</u>	
Species Found? <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No If not, why? _____	Reporter: <u>Richard A. Erickson</u>
Total No. Individuals <u>1</u> Subsequent Visit? <input type="checkbox"/> yes <input checked="" type="checkbox"/> no	Address: <u>LSA Associates 20 Executive Park</u>
Is this an existing NDDDB occurrence? <input type="checkbox"/> no <input checked="" type="checkbox"/> unk. Yes, Occ. # _____	<u>Suite 200, Irvine, CA 92614</u>
Collection? If yes: _____ Number _____ Museum / Herbarium _____	E-mail Address: <u>richard.erickson@LSA-assoc.com</u>
	Phone: <u>949 553-0666</u>

<b>Plant Information</b>	<b>Animal Information</b>
Phenology: _____% vegetative _____% flowering _____% fruiting	# adults _____ # juveniles _____ # larvae _____ # egg masses _____ # unknown _____
	<input type="checkbox"/> breeding <input type="checkbox"/> wintering <input type="checkbox"/> burrow site <input type="checkbox"/> rookery <input type="checkbox"/> nesting <input checked="" type="checkbox"/> other <u>territorial</u>

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

Puente Hills, Whittier Puente Hills Landfill Native

County: Los Angeles Landowner / Mgr.: Habitat Preservation Authority

Quad Name: Whittier and La Habra Elevation: 300-1000 ft

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  D Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  D GPS Make & Model \_\_\_\_\_

**DATUM:** NAD27  NAD83  WGS84  Horizontal Accuracy \_\_\_\_\_ meters/feet

Coordinate System: UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)

Coordinates: chat coordinates: ~33°58'18"N 117°59'50"W

**Habitat Description (plant communities, dominants, associates, substrates/soils, aspects/slope):**

coastal slope scrub, ruderal grassland, riparian scrub & woodland, eucalyptus woodland

Other rare taxa seen at THIS site on THIS date: Neotoma lepida intermedia  
(separate form preferred)

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor

Immediate AND surrounding land use: native habitat preserve

Visible disturbances: invasive exotic vegetation, extensive road system

Threats: oil extraction?

Comments: \_\_\_\_\_

<b>Determination:</b> (check one or more, and fill in blanks)	<b>Photographs:</b> (check one or more)	Slide	Print	Digital
<input type="checkbox"/> Keyed (cite reference): _____	Plant / animal	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/> Compared with specimen housed at: _____	Habitat	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/> Compared with photo / drawing in: _____	Diagnostic feature	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/> By another person (name): _____				
<input checked="" type="checkbox"/> Other: _____	May we obtain duplicates at our expense? yes <input type="checkbox"/> no <input type="checkbox"/>			

Mail to:

California Natural Diversity Database  
 Department of Fish and Game  
 1807 13<sup>th</sup> Street, Suite 202  
 Sacramento, CA 95811

Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov

For Office Use Only

Source Code \_\_\_\_\_ Quad Code \_\_\_\_\_  
 Elm Code \_\_\_\_\_ Occ. No. \_\_\_\_\_  
 EO Index No. \_\_\_\_\_ Map Index No. \_\_\_\_\_

Date of Field Work (mm/dd/yyyy): 6/13-7/9/2008

Reset

California Native Species Field Survey Form

Send Form

**Scientific Name:** Neotoma lepida intermedia

**Common Name:** San Diego desert woodrat

**Species Found?**  Yes  No If not, why?

Total No. Individuals ? Subsequent Visit?  yes  no

Is this an existing NDDDB occurrence?  no  unk. Yes, Occ. #

Collection? If yes: \_\_\_\_\_  
Number Museum / Herbarium

**Reporter:** Richard A. Erickson

**Address:** LSA Associates 20 Executive Park, Suite 200, Irvine, CA 92614

**E-mail Address:** richard.erickson@LSA-assoc.com

**Phone:** 949 553-0666

**Plant Information**

Phenology: \_\_\_\_\_% vegetative \_\_\_\_\_% flowering \_\_\_\_\_% fruiting

**Animal Information**

nest found

# adults	# juveniles	# larvae	# egg masses	# unknown
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
breeding	wintering	burrow site	rookery	nesting

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

Puente Hills, Whittier

County: Los Angeles Landowner / Mgr.: Puente Hills Landfill Native Habitat Preservation Authority

Quad Name: Whittier and La Habra Elevation: 300-1000 ft

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  D Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  D GPS Make & Model \_\_\_\_\_

**DATUM:** NAD27  NAD83  WGS84  Horizontal Accuracy \_\_\_\_\_ meters/feet

Coordinate System: UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)

Coordinates: woodrat nest coordinates: 33° 58' 17" N 118° 00' 20" W

**Habitat Description (plant communities, dominants, associates, substrates/soils, aspects/slope):**

coastal sage scrub, ruderal grassland, riparian scrub & woodland, eucalyptus woodland

Other rare taxa seen at THIS site on THIS date: Icterus virens  
 (separate form preferred)

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor

Immediate AND surrounding land use: native habitat preserve

Visible disturbances: invasive exotic vegetation, extensive road system

Threats: oil extraction?

Comments:

**Determination:** (check one or more, and fill in blanks)

Keyed (cite reference): \_\_\_\_\_

Compared with specimen housed at: \_\_\_\_\_

Compared with photo / drawing in: \_\_\_\_\_

By another person (name): \_\_\_\_\_

Other: \_\_\_\_\_

**Photographs:** (check one or more)

Slide	Print	Digital
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Plant / animal  
 Habitat  
 Diagnostic feature

May we obtain duplicates at our expense? yes  no



LSA ASSOCIATES, INC.  
PACIFIC CENTER  
703 PALOMAR AIRPORT RD., SUITE 260  
CARLSBAD, CALIFORNIA 92011

760.931.5471 TEL  
760.918.2458  
FAX

BERKELEY  
COLMA  
FORT COLLINS

IRVINE  
PALM SPRINGS  
POINT RICHMOND

RIVERSIDE  
ROCKLIN  
SAN LUIS OBISPO

August 4, 2009

Sandra Marquez  
U.S. Fish and Wildlife Service  
Carlsbad Field Office  
6010 Hidden Valley Road, Suite 101  
Carlsbad, California 92011

Lyann Comrack  
Nongame Wildlife Program  
California Department of Fish and Game  
1812 Ninth Street  
Sacramento, California 95811

Subject: Least Bell's Vireo and Coastal California Gnatcatcher Survey Results  
Puente Hills Landfill Native Habitat Preservation Authority Managed Lands  
City of Whittier, Los Angeles County, California (LSA Project No. PUE0901)

Dear Mss. Marquez and Comrack:

This letter report documents the results of protocol surveys for the least Bell's vireo (*Vireo bellii pusillus*) and coastal California gnatcatcher (*Polioptila californica californica*) conducted by LSA Associates, Inc. (LSA). Surveys were conducted within lands owned by the City of Whittier and managed by the Puente Hills Landfill Native Habitat Preservation Authority (Habitat Authority) in the City of Whittier, Los Angeles County, California (Figure 1: Project Location). The least Bell's vireo is a United States Fish and Wildlife Service (USFWS) federally endangered species and a California Department of Fish and Game (CDFG) endangered species. The coastal California gnatcatcher is a USFWS federally threatened species.

Protocol surveys of the Habitat Authority Lands detected no least Bell's vireos and a single coastal California gnatcatcher.

## STUDY AREA

The survey area is located within lands managed by the Habitat Authority. The survey area totals approximately 209 acres and is located within Sections 22, 23, 25, and 26 of Township 2 South, Range 11 West, as shown on the United States Geological Survey (USGS) 7.5-minute *Whittier* and *La Habra*, California quadrangles (Figure 1, attached).

The survey area is characterized by portions of the Arroyo Pescadero and La Cañada Verde drainages, adjacent hillsides, and access roads. Elevation ranges from approximately 300 to 1,000 feet above mean sea level. Vegetation communities within the survey area primarily include coastal sage scrub, riparian scrub, nonnative grassland, ruderal vegetation, eucalyptus woodland, ornamental vegetation, and other previously disturbed areas.



## METHODS

Mark J. Billings and Richard Erickson conducted nine protocol surveys for least Bell’s vireo and coastal California gnatcatcher from April 15 to June 29, 2009. During each of the surveys, the biologists walked slowly along the edge of and, when appropriate, through riparian and coastal sage scrub habitat, listening for least Bell’s vireos and coastal California gnatcatchers. Taped recordings of coastal California gnatcatchers were played periodically along the survey route during all of the surveys. With the aid of binoculars for viewing wildlife species, the surveying biologists waited for several minutes after each playing to look and listen both for least Bell’s vireos and for coastal California gnatcatchers.

**Table A: Survey Schedule and Conditions**

Date 2009	Time	Weather	Surveyors
April 15	0630–1030	Cool, mostly clear, and moderate wind	MJB, RE
April 28	0645–1000	Cool, partly overcast, and light wind	MJB, RE
May 8	0630–1000	Mild, clear, and calm	MJB, RE
May 18	0645–1000	Mild, mostly clear, and light wind	MJB, RE
May 28	0615–0900	Mild, overcast, and calm	RE
June 1	0600–0845	Mild, overcast, and calm	RE
June 8	0615–0930	Mild, overcast, and calm	MJB, RE
June 19	0630–0915	Mild, partly overcast, and calm	MJB, RE
June 29	0630–0945	Mild, mostly clear, and calm	MJB, RE

Surveyors: Mark J. Billings (MJB) and Richard Erickson (RE).

Surveys were conducted pursuant to Federal Fish and Wildlife Permit TE777965-8 (April 8, 2008–April 7, 2012) and a temporary authorization letter from the CDFG (May 12, 2003–March 31, 2007; renewal request submitted March 26, 2007 and approved on May 4, 2007, extending coverage indefinitely) in lieu of a Memorandum of Understanding between LSA and CDFG.

## RESULTS

No least Bell’s vireos were found during the surveys. On June 29, a single young male coastal California gnatcatcher was seen in the study area (Figure 2: 2008 and 2009 Survey Areas and Results, attached). The bird was most likely a recently fledged juvenile dispersing from its natal territory.

A complete list of the animals detected on these surveys is included in Table B (attached).

LSA ASSOCIATES, INC.

If you have any questions, please contact me at (760) 931-5471.

Sincerely,

LSA ASSOCIATES, INC.



Mark J. Billings  
Assistant Biologist

Attachments: Figure 1: Project Location  
Figure 2: 2008 and 2009 Survey Areas and Results  
Table B: Animal Species Observed  
California Native Species Field Survey Summary Form

**I CERTIFY THAT THE INFORMATION IN THIS SURVEY REPORT AND ATTACHED EXHIBITS FULLY AND ACCURATELY REPRESENTS MY WORK:**

**SURVEYOR:**

**PERMIT NUMBER**


**DATE:**



TE-777965-8

August 4, 2009

Mark J. Billings



TE-777965-8

August 4, 2009

Richard Erickson







FIGURE 1

LSA

LEGEND


- 2009 Survey Area (approx. 209 acres)
- 2008 Survey Area (approx. 111 acres)





Puente Hills Landfill Native Habitat Preservation Authority  
 City of Whittier Oil Exploration  
 Project Location

SOURCE: USGS 7.5' QUAD - WHITTIER (81) AND LA HABRA (81); CALIF.





LSA



SOURCE: DigitalGlobe (April 2008)

I:\PUE0901\GIS\Reports\Bird\_Survey\Fig2\_Survey\_Results.mxd (07/15/2009)

- Coastal California Gnatcatcher Observed June 29, 2009
  - Coastal California Gnatcatcher Observed June 9 & 10, 2008
  - Most Suitable Coastal California Gnatcatcher Habitat
  - Potential Least Bell's Vireo Habitat
- 2009 Survey Area
  - 2008 Survey Area

FIGURE 2

Puente Hills Landfill Native Habitat Preservation Authority  
 City of Whittier Oil Exploration  
 2008 and 2009 Survey Areas and Results

## TABLE B

### ANIMAL SPECIES OBSERVED

This is a list of the conspicuous aerial insects, amphibians (none observed), reptiles, birds, and mammals noted in the study area by LSA biologists. Presence may be noted if a species is seen or heard, or identified by the presence of tracks, scat, or other signs.

- \* Species not native to the study area
- \*\* Special-status species (note that some species are only considered special-status if nesting colonies are present)

Scientific Name	Common Name
<b>ANISOPTERA</b>	<b>TYPICAL DRAGONFLIES</b>
<b>Libellulidae</b>	<b>Cruisers, Emeralds, Baskettails, and Skimmers</b>
<i>Libellula saturata</i>	Flame skimmer
<i>Pantala flavescens</i>	Wandering glider
<i>Pantala hymenaea</i>	Spot-winged glider
<b>LEPIDOPTERA</b>	<b>BUTTERFLIES</b>
<b>Papilionidae</b>	<b>Swallowtails</b>
<i>Papilio rutulus</i>	Western tiger swallowtail
<b>Pieridae</b>	<b>Whites and Sulphurs</b>
<i>Pontia protodice</i>	Checkered white
* <i>Pieris rapae</i>	Cabbage white
<i>Anthocharis sara</i>	Sara orangetip
<b>Lycaenidae</b>	<b>Gossamer-Wing Butterflies</b>
<i>Leptotes marina</i>	Marine blue
<i>Plebejus acmon</i>	Acmon blue
<b>Nymphalidae</b>	<b>Brush-Footed Butterflies</b>
<i>Agraulis vanillae</i>	Gulf fritillary
<i>Nymphalis antiopa</i>	Mourning cloak
<i>Vanessa cardui</i>	Painted lady
<i>Vanessa atalanta</i>	Red admiral
<i>Danaus plexippus</i>	Monarch
<b>REPTILIA</b>	<b>REPTILES</b>
<b>Phrynosomatidae</b>	<b>Phrynosomatid Lizards</b>
<i>Sceloporus occidentalis</i>	Western fence lizard
<b>AVES</b>	<b>BIRDS</b>
<b>Anatidae</b>	<b>Ducks, Geese, and Swans</b>
<i>Anas platyrhynchos</i>	Mallard
<b>Odontophoridae</b>	<b>New World Quail</b>
<i>Callipepla californica</i>	California quail
<b>Phalacrocoracidae</b>	<b>Cormorants</b>
** <i>Phalacrocorax auritus</i>	Double-crested cormorant

<b>Ardeidae</b>	<b>Hérons, Bitterns, and Allies</b>
** <i>Ardea herodias</i>	Great blue heron
** <i>Ardea alba</i>	Great egret
<b>Cathartidae</b>	<b>New World Vultures</b>
<i>Cathartes aura</i>	Turkey vulture
<b>Accipitridae</b>	<b>Hawks, Kites, Eagles, and Allies</b>
** <i>Accipiter striatus</i>	Sharp-shinned hawk
** <i>Accipiter cooperii</i>	Cooper's hawk
<i>Buteo lineatus</i>	Red-shouldered hawk
<i>Buteo jamaicensis</i>	Red-tailed hawk
<b>Falconidae</b>	<b>Caracaras and Falcons</b>
<i>Falco sparverius</i>	American kestrel
<b>Scolopacidae</b>	<b>Sandpipers, Phalaropes, and Allies</b>
<i>Numenius phaeopus</i>	Whimbrel
<b>Laridae</b>	<b>Gulls, Terns, and Skimmers</b>
<i>Larus occidentalis</i>	Western gull
<i>Larus californicus</i>	California gull
<i>Larus glaucescens</i>	Glaucous-winged gull
** <i>Hydroprogne caspia</i>	Caspian tern
<b>Columbidae</b>	<b>Pigeons and Doves</b>
* <i>Columba livia</i>	Rock (Feral) pigeon
<i>Patagioenas fasciata</i>	Band-tailed pigeon
<i>Zenaida macroura</i>	Mourning dove
<b>Psittacidae</b>	<b>Lories, Parakeets, Macaws, and Parrots</b>
* <i>Amazona viridigenalis</i>	Red-crowned parrot
<b>Cuculidae</b>	<b>Cuckoos, Roadrunners, and Anis</b>
<i>Geococcyx californianus</i>	Greater roadrunner
<b>Apodidae</b>	<b>Swifts</b>
** <i>Chaetura vauxi</i>	Vaux's swift
<i>Aeronautes saxatilis</i>	White-throated swift
<b>Trochilidae</b>	<b>Hummingbirds</b>
<i>Archilochus alexandri</i>	Black-chinned hummingbird
** <i>Calypte anna</i>	Anna's hummingbird
** <i>Selasphorus sasin</i>	Allen's hummingbird
<b>Picidae</b>	<b>Woodpeckers and Allies</b>
<i>Melanerpes formicivorus</i>	Acorn woodpecker
** <i>Picoides nuttallii</i>	Nuttall's woodpecker
<i>Picoides pubescens</i>	Downy woodpecker
<i>Colaptes auratus</i>	Northern flicker
<b>Tyrannidae</b>	<b>Tyrant Flycatchers</b>
<i>Contopus sordidulus</i>	Western wood-pewee
<i>Empidonax difficilis</i>	Pacific-slope flycatcher
<i>Sayornis nigricans</i>	Black phoebe
<i>Myiarchus cinerascens</i>	Ash-throated flycatcher
<i>Tyrannus vociferans</i>	Cassin's kingbird
<b>Vireonidae</b>	<b>Vireos</b>
<i>Vireo huttoni</i>	Hutton's vireo
<i>Vireo gilvus</i>	Warbling vireo
<b>Corvidae</b>	<b>Crows and Jays</b>
<i>Aphelocoma californica</i>	Western scrub-jay



<i>Corvus brachyrhynchos</i>	American crow
<i>Corvus corax</i>	Common raven
<b>Alaudidae</b>	<b>Larks</b>
** <i>Eremophila alpestris</i>	Horned lark
<b>Hirundinidae</b>	<b>Swallows</b>
<i>Stelgidopteryx serripennis</i>	Northern rough-winged swallow
<i>Petrochelidon pyrrhonota</i>	Cliff swallow
<i>Hirundo rustica</i>	Barn swallow
<b>Aegithalidae</b>	<b>Long-Tailed Tits and Bushtits</b>
<i>Psaltriparus minimus</i>	Bushtit
<b>Troglodytidae</b>	<b>Wrens</b>
<i>Thryomanes bewickii</i>	Bewick's wren
<i>Troglodytes aedon</i>	House wren
<b>Sylviidae</b>	<b>Old World Warblers and Gnatcatchers</b>
<i>Polioptila caerulea</i>	Blue-gray gnatcatcher
** <i>Polioptila californica californica</i>	Coastal California gnatcatcher
<b>Turdidae</b>	<b>Thrushes</b>
<i>Sialia mexicana</i>	Western bluebird
<i>Catharus ustulatus</i>	Swainson's thrush
<i>Catharus guttatus</i>	Hermit thrush
<i>Turdus migratorius</i>	American robin
<b>Timaliidae</b>	<b>Babblers</b>
<i>Chamaea fasciata</i>	Wrentit
<b>Mimidae</b>	<b>Mockingbirds and Thrashers</b>
<i>Mimus polyglottos</i>	Northern mockingbird
<i>Toxostoma redivivum</i>	California thrasher
<b>Bombycillidae</b>	<b>Waxwings</b>
<i>Bombycilla cedrorum</i>	Cedar waxwing
<b>Ptilonotidae</b>	<b>Silky-flycatchers</b>
<i>Phainopepla nitens</i>	Phainopepla
<b>Parulidae</b>	<b>Wood Warblers</b>
<i>Vermivora celata</i>	Orange-crowned warbler
<i>Vermivora ruficapilla</i>	Nashville warbler
** <i>Dendroica petechia</i>	Yellow warbler
<i>Dendroica coronata</i>	Yellow-rumped warbler
<i>Dendroica nigrescens</i>	Black-throated gray warbler
<i>Dendroica townsendi</i>	Townsend's warbler
<i>Oporornis tolmiei</i>	MacGillivray's warbler
<i>Geothlypis trichas</i>	Common yellowthroat
<i>Wilsonia pusilla</i>	Wilson's warbler
** <i>Icteria virens</i>	Yellow-breasted chat
<b>Thraupidae</b>	<b>Tanagers</b>
<i>Piranga ludoviciana</i>	Western tanager
<b>Emberizidae</b>	<b>Emberizids</b>
<i>Pipilo maculatus</i>	Spotted towhee
<i>Pipilo crissalis</i>	California towhee
** <i>Aimophila ruficeps</i>	Rufous-crowned sparrow
** <i>Spizella breweri</i>	Brewer's sparrow
<i>Passerella iliaca</i>	Fox sparrow
<i>Melospiza melodia</i>	Song sparrow

<i>Zonotrichia leucophrys</i>	White-crowned sparrow
<i>Zonotrichia atricapilla</i>	Golden-crowned sparrow
<b>Cardinalidae</b>	<b>Cardinals, Saltators, and Allies</b>
<i>Pheucticus ludovicianus</i>	Rose-breasted grosbeak
<i>Pheucticus melanocephalus</i>	Black-headed grosbeak
<i>Passerina caerulea</i>	Blue grosbeak
<i>Passerina amoena</i>	Lazuli bunting
<b>Icteridae</b>	<b>Blackbirds</b>
<i>Euphagus cyanocephalus</i>	Brewer's blackbird
<i>Icterus cucullatus</i>	Hooded oriole
<i>Icterus bullockii</i>	Bullock's oriole
<b>Fringillidae</b>	<b>Fringilline and Cardueline Finches and Allies</b>
<i>Carpodacus mexicanus</i>	House finch
<i>Carduelis psaltria</i>	Lesser goldfinch
<i>Carduelis tristis</i>	American goldfinch
<b>Passeridae</b>	<b>Old World Sparrows</b>
* <i>Passer domesticus</i>	House sparrow
<b>Estrildidae</b>	<b>Estrildid Finches</b>
* <i>Lonchura punctulata</i>	Nutmeg mannikin
<b>MAMMALIA</b>	<b>MAMMALS</b>
<b>Sciuridae</b>	<b>Squirrels, Chipmunks, and Marmots</b>
<i>Spermophilus beecheyi</i>	California ground squirrel
<b>Geomyidae</b>	<b>Pocket Gophers</b>
<i>Thomomys bottae</i>	Botta's pocket gopher
<b>Cricetidae</b>	<b>Hamsters, Voles, Lemmings, and New World Rats and Mice</b>
** <i>Neotoma lepida</i>	San Diego desert woodrat
<i>Neotoma macrotis</i>	Big-eared woodrat
<b>Leporidae</b>	<b>Rabbits and Hares</b>
<i>Sylvilagus audubonii</i>	Audubon's cottontail
<b>Felidae</b>	<b>Cats</b>
<i>Lynx rufus</i>	Bobcat
<b>Canidae</b>	<b>Foxes, Wolves, and Allies</b>
<i>Canis latrans</i>	Coyote
<b>Cervidae</b>	<b>Deer, Elk, and Allies</b>
<i>Odocoileus hemionus</i>	Mule deer

**Taxonomy and nomenclature are based on the following.**

**Damselflies and dragonflies:** Manolis, T. (2003, *Dragonflies and Damselflies of California*, University of California Press, Berkeley).

**Butterflies:** North American Butterfly Association (2001, *NABA checklist and English Names of North American Butterflies*, Second Edition, North American Butterfly Association, Morristown, New Jersey).

**Amphibians and reptiles:** Crother, B.I. ed. (2008. *Scientific and Standard English Names of Amphibians and Reptiles of North America North of Mexico. Herpetological Circular 37*) for species

taxonomy and nomenclature; Stebbins, R.C. (2003, A Field Guide to Western Reptiles and Amphibians, third edition, Houghton Mifflin, Boston) for sequence and higher order taxonomy.

**Birds:** American Ornithologists' Union (1998, The A.O.U. Checklist of North American Birds, Seventh Edition, American Ornithologists' Union, Washington D.C.; and supplements; see <http://aou.org.whsites.net/checklist/index.php3>).

**Mammals:** Wilson, D.E., and D.M. Reeder, eds. (2005. Mammal Species of the World, 3rd ed. Johns Hopkins University Press, Baltimore, Maryland; see <http://nmnhgoph.si.edu/msw/>).



Mail to:  
 California Natural Diversity Database  
 Department of Fish and Game  
 1807 13<sup>th</sup> Street, Suite 202  
 Sacramento, CA 95811

Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov

*For Office Use Only*

Source Code \_\_\_\_\_ Quad Code \_\_\_\_\_  
 Elm Code \_\_\_\_\_ Occ. No. \_\_\_\_\_  
 EO Index No. \_\_\_\_\_ Map Index No. \_\_\_\_\_

Date of Field Work (mm/dd/yyyy): 6/29/09

**Reset**

**California Native Species Field Survey Form**

**Send Form**

Scientific Name: Vireo bellii pusillus

Common Name: least Bell's vireo

Species Found?  Yes  No not present  
If not, why?

Total No. Individuals \_\_\_\_\_ Subsequent Visit?  yes  no  
 Is this an existing NDDB occurrence?  no  unk.  
Yes, Occ. #

Collection? If yes: \_\_\_\_\_  
Number Museum / Herbarium

Reporter: Mark J. Billings  
 Address: 703 Palomar Airport Road, Suite 260  
Carlsbad, California 92011  
 E-mail Address: mark.billings@lsa-assoc.com  
 Phone: (760) 931-5471

**Plant Information**

Phenology: \_\_\_\_\_% vegetative \_\_\_\_\_% flowering \_\_\_\_\_% fruiting

**Animal Information**

# adults # juveniles # larvae # egg masses # unknown  
 breeding  wintering  burrow site  rookery  nesting  other

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

County: Los Angeles Landowner / Mgr.: \_\_\_\_\_  
 Quad Name: \_\_\_\_\_ Elevation: \_\_\_\_\_  
 T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian: H  M  S  Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_  
 T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian: H  M  S  GPS Make & Model \_\_\_\_\_  
**DATUM:** NAD27  NAD83  WGS84  Horizontal Accuracy \_\_\_\_\_ meters/feet  
**Coordinate System:** UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)   
 Coordinates: 33°57'42.54" N, 118°00'07.10" W

**Habitat Description (plant communities, dominants, associates, substrates/soils, aspects/slope):**  
coastal sage scrub, riparian scrub, nonnative grassland, ruderal vegetation, eucalyptus woodland, and ornamental vegetation

Other rare taxa seen at THIS site on THIS date:  
 (separate form preferred)

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor

Immediate AND surrounding land use:  
 Visible disturbances: development  
 Threats: development  
 Comments:

**Determination:** (check one or more, and fill in blanks)

Keyed (cite reference): \_\_\_\_\_  
 Compared with specimen housed at: \_\_\_\_\_  
 Compared with photo / drawing in: \_\_\_\_\_  
 By another person (name): \_\_\_\_\_  
 Other: \_\_\_\_\_

**Photographs:** (check one or more)

Slide	Print	Digital
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

May we obtain duplicates at our expense? yes  no

Mail to:  
 California Natural Diversity Database  
 Department of Fish and Game  
 1807 13<sup>th</sup> Street, Suite 202  
 Sacramento, CA 95811

Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov

For Office Use Only	
Source Code _____	Quad Code _____
Elm Code _____	Occ. No. _____
EO Index No. _____	Map Index No. _____

Date of Field Work (mm/dd/yyyy): 6/29/09

**Reset**

### California Native Species Field Survey Form

**Send Form**

Scientific Name: Poliioptila californica californica

Common Name: coastal California gnatcatcher

Species Found?  Yes  No  
 If not, why? \_\_\_\_\_  
 Total No. Individuals 1 Subsequent Visit?  yes  no  
 Is this an existing NDDDB occurrence?  no  unk.  
 Yes, Occ. # \_\_\_\_\_  
 Collection? If yes: \_\_\_\_\_  
 Number \_\_\_\_\_ Museum / Herbarium \_\_\_\_\_

Reporter: Mark J. Billings for Richard Erickson  
 Address: 703 Palomar Airport Road, Suite 260  
Carlsbad, California 92011  
 E-mail Address: mark.billings@lsa-assoc.com  
 Phone: (760) 931-5471

**Plant Information**  
 Phenology: \_\_\_\_\_% vegetative \_\_\_\_\_% flowering \_\_\_\_\_% fruiting

**Animal Information**  
1  
 # adults \_\_\_\_\_ # juveniles \_\_\_\_\_ # larvae \_\_\_\_\_ # egg masses \_\_\_\_\_ # unknown \_\_\_\_\_  
 breeding  wintering  burrow site  rookery  nesting  other

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

County: Los Angeles Landowner / Mgr.: \_\_\_\_\_  
 Quad Name: \_\_\_\_\_ Elevation: \_\_\_\_\_  
 T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_  
 T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4, Meridian:  H  M  S  GPS Make & Model \_\_\_\_\_  
**DATUM:** NAD27  NAD83  WGS84  Horizontal Accuracy \_\_\_\_\_ meters/feet  
**Coordinate System:** UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)   
**Coordinates:** 33°57'42.54" N, 118°00'07.10" W

**Habitat Description (plant communities, dominants, associates, substrates/soils, aspects/slope):**  
coastal sage scrub, riparian scrub, nonnative grassland, ruderal vegetation, eucalyptus woodland, and ornamental vegetation

Other rare taxa seen at THIS site on THIS date:  
 (separate form preferred)

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor  
 Immediate AND surrounding land use:  
 Visible disturbances: development  
 Threats: development  
 Comments:

**Determination:** (check one or more, and fill in blanks)  
 Keyed (cite reference): \_\_\_\_\_  
 Compared with specimen housed at: \_\_\_\_\_  
 Compared with photo / drawing in: \_\_\_\_\_  
 By another person (name): \_\_\_\_\_  
 Other: \_\_\_\_\_

**Photographs:** (check one or more) Slide Print Digital  
 Plant / animal     
 Habitat     
 Diagnostic feature     
 May we obtain duplicates at our expense? yes  no

Mail to:  
 California Natural Diversity Database  
 Department of Fish and Game  
 1807 13<sup>th</sup> Street, Suite 202  
 Sacramento, CA 95811

Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov

*For Office Use Only*

Source Code \_\_\_\_\_ Quad Code \_\_\_\_\_  
 Elm Code \_\_\_\_\_ Occ. No. \_\_\_\_\_  
 EO Index No. \_\_\_\_\_ Map Index No. \_\_\_\_\_

Date of Field Work (mm/dd/yyyy): 6/29/09

**Reset**

## California Native Species Field Survey Form

**Send Form**

**Scientific Name:** Icteria virens

**Common Name:** yellow-breasted chat

**Species Found?**  Yes  No If not, why?

Total No. Individuals 1 Subsequent Visit?  yes  no

Is this an existing NDDDB occurrence?  no  unk.

Collection? If yes: \_\_\_\_\_  
Number Museum / Herbarium

**Reporter:** Mark J. Billings for Richard Erickson

**Address:** 703 Palomar Airport Road, Suite 260  
Carlsbad, California 92011

**E-mail Address:** mark.billings@lsa-assoc.com

**Phone:** (760) 931-5471

**Plant Information**

Phenology: \_\_\_\_\_% vegetative \_\_\_\_\_% flowering \_\_\_\_\_% fruiting

**Animal Information**

1 # adults # juveniles # larvae # egg masses # unknown

breeding  wintering  burrow site  rookery  nesting  other

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

County: Los Angeles Landowner / Mgr.: \_\_\_\_\_

Quad Name: \_\_\_\_\_ Elevation: \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ ¼ of \_\_\_\_\_ ¼, Meridian:  H  M  S  Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_

T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ ¼ of \_\_\_\_\_ ¼, Meridian:  H  M  S  GPS Make & Model \_\_\_\_\_

**DATUM:**  NAD27  NAD83  WGS84  Horizontal Accuracy \_\_\_\_\_ meters/feet

**Coordinate System:** UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)

**Coordinates:** 33°57'42.54" N, 118°00'07.10" W

**Habitat Description (plant communities, dominants, associates, substrates/soils, aspects/slope):**

coastal sage scrub, riparian scrub, nonnative grassland, ruderal vegetation, eucalyptus woodland, and ornamental vegetation

Other rare taxa seen at THIS site on THIS date: (separate form preferred)

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor

Immediate AND surrounding land use:

Visible disturbances: development

Threats: development

Comments:

**Determination:** (check one or more, and fill in blanks)

Keyed (cite reference): \_\_\_\_\_

Compared with specimen housed at: \_\_\_\_\_

Compared with photo / drawing in: \_\_\_\_\_

By another person (name): \_\_\_\_\_

Other: \_\_\_\_\_

**Photographs:** (check one or more)

Plant / animal  Slide  Print  Digital

Habitat  Slide  Print  Digital

Diagnostic feature  Slide  Print  Digital

May we obtain duplicates at our expense? yes  no



Mail to:  
 California Natural Diversity Database  
 Department of Fish and Game  
 1807 13<sup>th</sup> Street, Suite 202  
 Sacramento, CA 95811

Fax: (916) 324-0475 email: CNDDDB@dfg.ca.gov

For Office Use Only	
Source Code _____	Quad Code _____
Elm Code _____	Occ. No. _____
EO Index No. _____	Map Index No. _____

Date of Field Work (mm/dd/yyyy): 6/19/09

**Reset**

### California Native Species Field Survey Form

**Send Form**

Scientific Name: Neotoma lepida

Common Name: desert woodrat

Species Found?  Yes  No If not, why?  
 Total No. Individuals \_\_\_\_\_ Subsequent Visit?  yes  no  
 Is this an existing NDDDB occurrence? \_\_\_\_\_  no  unk.  
Yes, Occ. #  
 Collection? If yes: \_\_\_\_\_  
Number Museum / Herbarium

Reporter: Mark J. Billings for Richard Erickson  
 Address: 703 Palomar Airport Road, Suite 260  
Carlsbad, California 92011  
 E-mail Address: mark.billings@lsa-assoc.com  
 Phone: (760) 931-5971

**Plant Information**  
 Phenology: \_\_\_\_\_% vegetative \_\_\_\_\_% flowering \_\_\_\_\_% fruiting

**Animal Information**  
 # adults # juveniles # larvae # egg masses # unknown  
 breeding  wintering  burrow site  rookery  nesting  other

**Location Description (please attach map AND/OR fill out your choice of coordinates, below)**

County: Los Angeles Landowner / Mgr.: \_\_\_\_\_  
 Quad Name: \_\_\_\_\_ Elevation: \_\_\_\_\_  
 T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ ¼ of \_\_\_\_\_ ¼, Meridian: H  M  S  Source of Coordinates (GPS, topo. map & type): \_\_\_\_\_  
 T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_, \_\_\_\_\_ ¼ of \_\_\_\_\_ ¼, Meridian: H  M  S  GPS Make & Model \_\_\_\_\_  
**DATUM:** NAD27  NAD83  WGS84  Horizontal Accuracy \_\_\_\_\_ meters/feet  
**Coordinate System:** UTM Zone 10  UTM Zone 11  OR Geographic (Latitude & Longitude)   
**Coordinates:** 33°57'42.54" N, 118°00'07.10" W

**Habitat Description (plant communities, dominants, associates, substrates/soils, aspects/slope):**  
coastal sage scrub, riparian scrub, nonnative grassland, ruderal vegetation, eucalyptus woodland, and ornamental vegetation

Other rare taxa seen at THIS site on THIS date:  
 (separate form preferred)

**Site Information** Overall site/occurrence quality/viability (site + population):  Excellent  Good  Fair  Poor  
 Immediate AND surrounding land use:  
 Visible disturbances: development  
 Threats: development  
 Comments:

**Determination:** (check one or more, and fill in blanks)  
 Keyed (cite reference): \_\_\_\_\_  
 Compared with specimen housed at: \_\_\_\_\_  
 Compared with photo / drawing in: \_\_\_\_\_  
 By another person (name): \_\_\_\_\_  
 Other: \_\_\_\_\_

**Photographs:** (check one or more) Slide Print Digital  
 Plant / animal     
 Habitat     
 Diagnostic feature     
 May we obtain duplicates at our expense? yes  no